



Medicinal Plants as Aphrodisiac Agents: A Current Status

Saurabh Nimesh^{1*}, Shubham², Vrish Dhvaj Ashwlayan³ and Pankaj Barman⁴

¹Department of Pharmaceutical Technology, Meerut Institute of Engineering and Technology, Meerut, Uttar Pradesh, India

²Senior Veterinary Field Manager; Vet Mankind Pharma Ltd., Haldwani Headquarter, Uttarakhand, India

³Associate Professor; Department of Pharmaceutical Technology, Meerut Institute of Engineering and Technology, Meerut, Uttar Pradesh, India

⁴M. Pharm. (Pharmacology) Student; Department of Pharmaceutical Technology, Meerut Institute of Engineering and Technology, Meerut, Uttar Pradesh, India

*Corresponding Author: Saurabh Nimesh, Department of Pharmaceutical Technology, Meerut Institute of Engineering and Technology, Meerut, Uttar Pradesh, India.

Received: May 21, 2019; Published: July 25, 2019

DOI: 10.31080/ASPS.2019.03.0355

Abstract

Modern life vogue and bound environmental exposures have resulted in male infertility. The activating factors turn out differing types of derangements that directly or indirectly cause sexual dysfunctions. Male impotence conjointly known as erectile dysfunction (ED) may be a common medical condition that affects the sexual lifetime of ample men worldwide. ED is outlined because the inability of a person to realize and maintain an erection adequate for naturally satisfactory intercourse. This literary criticism discusses regarding aphrodisiac potential of plants, its biological science name, Common name, family, parts used and chemical constituents, that are useful for investigator to development new aphrodisiac formulations. Hence, patients are seeking complementary and practice of medicine to treat sexual dysfunction. Ayurveda and different Indian literature mention the utilization of plants in numerous human ailments. India has regarding over 45000 plant species and among them many thousand are claimed to possess medicative properties.

Keywords: Sildenafil; Ayurveda; India; chemistry; Aphrodite; Fabaceae

Abbreviations

WHO: World Health Organization; ED: Erectile Dysfunction; NO: Nitric Oxide; ML: Mount Latency; IL: Intromission Latency; EL: Ejaculation Latency; MF: Mounting Frequency; IF: Intromission Frequency; PEI: Post-Ejaculatory Interval; Pgs: Phosphodiesterase; FSH: Follicle Stimulating Hormone

Introduction

Aphrodisiac is that the word derived from Aphrodite, the Greek god of sexual, love and sweetness. Associate aphrodisiac is outlined as an agent (food or drug) that arouses physical attraction or sexual desire [1]. The chance of bioactive aphrodisiacs which can be derived from plants, animals or minerals, has been engaging throughout recorded history [2]. Aphrodisiac are mentioned there as Vajikaranas, the word vaji that means horse and karanta meaning creating i.e. Live to excite lust by charms etc. Natural products are on the market in texts of Ayurveda for their spermatogenic and virility potential activities. Ayurvedic aphrodisiac medical specialty is classified into vajikarana (pharmacological) and rasayana (non-pharmacological products) [3]. The plant-based, ancient or

traditional medicine systems still play an important role in health care, with regarding 80% of the world's inhabitants relying in the main on ancient medicines for his or her primary health care. Modern pharmacopoeia still contains a minimum of 25% drugs derived from plants and plenty of others, that are artificial or synthetic analogues, designed on model compounds isolated from plants. Medicinal herbal plants produce bioactive compounds used in the main for medicinal functions [4-6]. Some well-known herbal aphrodisiacs are genus *Allium sativum*, *Alpinia galangal*, *Anacardium occidentale*, *Anacyclus pyrethrum*, *Butea frondosa*, *Caesalpinia benthamiana*, *Cannabis sativa*, *Chlorophyllum borivilianum*, *Citrullus lanatus*, *Eurycoma longifolia*, *Ginkgo biloba*, *Hibiscus sabdariffa*, etc. Sexual relationships are a some of the foremost necessary social and biological relationship in human life. According to World Health Organization (WHO) Sexual health is prime to the physical or emotional health and wellbeing of people, couples and families and to the social or economic development of communities and countries [7,8]. The National Institutes of Health Consensus Development Conference on Impotence (7 December 1992) has outlined, Male impotence conjointly known as ED may be a common medical

condition that affects the sexual life of millions of men worldwide. Impotency or ED as the 'inability to realize and maintain a penial erection adequate for satisfactory sexual relationship' (Figure 1) [9]. ED is outlined because the persistent inability to get associated maintain an erection comfortable for naturally satisfactory intercourse. Male reproductive capability was found to be deficient in nearly 50% of infertile couples in step with a study carried by the WHO. Sexual disfunction may be a serious medical and social symptom that happens in 10 - 52% of men and 25 - 63% of women [10-12].




Figure 1: Spectrum of Erectile dysfunction.

Mechanism involved in aphrodisiac potentials (Table 1)

Sexual desire is controlled and regulated by the central nervous system that integrates tactile, olfactive and mental stimuli (Figure 2) [13].




Figure 2: Mechanism of erection.

Side effects of Allopathic treatments used in sexual dysfunction

Side effects include drowsiness, insomnia, nasal congestion, headaches, dizziness, tachycardia, weight loss, etc. (Figure 3) [19].

S. No.	Stages	Explanation
1	First	Some aphrodisiac merely provides a burst of nutritious worth rising the immediate health or well-being of the patron and consequently improving sexual performance and concupiscence (libido).
2	Second	This cluster includes the supposed aphrodisiac have a lot of specific physiological affects however don't seem to be psychologically active. They will have an effect on blood flow; increase duration of sexual intercourse by desensitizing the sex organ space [14,15].
3	Third	The third cluster of aphrodisiac is created up compounds that are psychopharmacological, i.e. they really cross the blood brain barriers and stimulates some space of arousal [16]. This class includes a wide range of neurotransmitters, hormones, pheromones and drugs that interfere with the traditional perform of those molecules [17]. This class is most tough to check as a result of information of each arousal and therefore the mechanisms of the psychoactive properties of drugs are restricted. Solely the foremost general data regarding arousal and therefore the brain is known [18].

Table 1




Figure 3: Side effects of sildenafil, avanafil, tadalafil and vardenafil.

Some medicinal plants with aphrodisiac potential

Some of the traditional plants have tested to possess a conventional similarly as scientifically proven aphrodisiac which will enhance passion, increase physical attraction, enhance sexual performance and facilitate to extend the intensity of sexual love [20]. A short report of aphrodisiac plants in table 2 [21-26].

Scientific name (Family)	Common name	Pharmacology	Mechanism of action	Chemistry	Class of isolates
<i>Allium sativum</i> (Amaryllidaceae)	Garlic	The alcoholic extract of <i>A. sativum</i> increased sexual behaviour through the activities of sulphated compounds, peptides, flavonoids and phenolics	Allicin increases blood flow to sexual organs through nitric oxide (NO) synthase	Peptides, sulphated compounds, steroids, flavonoids, volatile oils with sulphated compounds like alliin, enzymes, minerals and vitamins	Peptides, steroids, terpenes, flavonoids, volatile oils and vitamins
<i>Alpinia galanga</i> (Zingiberaceae)	Greater galangal, blue ginger	Methanolic extract of <i>A. galangal</i> showed increase in serum testosterone levels at 300 mg/kg/day		Spectroscopic analysis of sample has revealed the presence of 1'S'-1'-acetoxychavicol acetate, 1'S'-1'-acetoxyeugenol acetate, 1'S'-1' hydroxychavicol acetate, trans-p-hydroxycinnam-aldehyde, trans-p-coumaryl alcohol, trans-p hydroxycinnamyl acetate, β - bisabolone and β -selinene	Coumarin, terpenoids, flavonoids, volatile oils, and phenols
<i>Anacardium occidentale</i> (Anacardiaceae)	Cashew	In a study to determine the aphrodisiac activity of the oils from <i>A. occidentale</i> seeds and shell, the result showed significant increase in sexual parameters		2-hydroxy-6-pentadecylbenzoic acid, the ethanolic extract of the nuts of <i>A. occidentale</i> contains phytochemicals such as phenols, carbohydrates, proteins and xanthoproteins as well as volatile oils, 2,6-dihydroxybenzoic acid from cashew apple, myristicin, kaempferol, rhamnetin, cyanidin, peonidin, delphinidin which are flavonoid compounds. Other isolated compounds are 2- hydroxy-6-pentadecylbenzoic acid, cardinal and salicylic acid	Carbohydrates, phenols, flavonoids, steroids, and proteins
<i>Anacyclus pyrethrum</i> (Asteraceae)	Arkakara	Administration of 50 mg/kg and 100 mg/kg of aqueous extract in albino rats showed significant anabolic and spermatogenic effects. In a separate study, petroleum ether extract had marked influence on body weight and accessory sexual organs weight as compared with arachis oil	This could be partly explained by its vasorelaxant properties which may be caused by an increase in NO production in vascular bed and a decrease in its destruction	Alkyl amides, pyrethrins, inulin, sesamine, hydrocaroline, pellitorine, volatile oils such as it is also composed of 2-phenyl ethylamine, anacylin, β -biotol, salvia-4(14)-en-1-one. Eudesma-4(15),7- diene-1-ol and β -himachalol; the essential oil also contains germacrema D, germacrema-4(15),5,10(14) trien-1-a-ol, caryophyllene oxide, cedryl acetate, eudesma-4(15),7- diene-1- β -ol and spathuleno	Amides, and Volatile oils
<i>Butea frondosa</i> (Papilionaceae)	Flame of the forest, bastard teak	The extract (400 mg/kg body wt./day) was administered orally by gavage for 28 days. Mount latency (ML), intromission latency (IL), ejaculation latency (EL), mounting frequency (MF), intromission frequency (IF), ejaculation frequency (EF) and post-ejaculatory interval (PEI) were the parameters observed before and during the sexual behaviour study at day 0, 7, 10, 14, 21, and 28. The extract reduced significantly ML, IL, EL and PEI ($p < 0.05$). The extract also increased significantly MF, IF and EF ($p < 0.05$). These effects were observed in sexually active and inactive male rats		Fixed oil 18%, Water soluble albuminoid substances 19% and glucose 6%. Fatty acids isolated from this oil are orleic linoleic, lenorlenic, palmitic, stearic, arachidic, behenic and lingo cleric acid. Q-hydroxy-1-methyo allophonic acid, 15-hydroxy pentasonic acid and 1carboxy methoxy-2-carboxy hydrazine have been isolated from the seed coat. Seed has shown the presence of alkaloid monsperrmine from the alcoholic extract of the seeds are identified palasonin and palasonin-N-Phenyl imidine. Aqueous methanolic extract contains a triazine compound, 4-arbomethoxy-3-dioxo-hydro-1,2,4-triazine 4.Carboxymethoxy 3.6 dioxo-hydro 1, 2, 4, triazine	Amino acids, alkaloids, and fixed oils

<i>Caesalpinia benthamiana</i> (<i>Caesalpinia-aceae</i>)	Bail	The methanolic extract exhibited an accelerator effect by decreasing the latent time. The oral administration of aqueous extract of <i>C. benthamiana</i> showed significant increase in mounting frequency and intromission frequency the dosage of 50 mg/kg		The petroleum ether extract of the bark has yielded cassane diterpenes with antibacterial activity such as deoxycassaldehyde C, benthaminine I and benthaminine 2, the aqueous extract contains flavonoids, phenols, anthraquinones such as gallic acid, esveratrol; the chloroform and n-butanol extract contains methyl gallate, shikimic acid-3-O-gallate, 1-O-methyl-D-chiroinositol, (-)-epicatechin	Terpenes, benthamine, fatty acids, flavonoids, and alkaloids
<i>Cannabis sativa</i> (<i>Cannabina-ceae</i>)	Marijuana, bhaang	In India's Ayurveda and Chinese, Unani medicine, cannabis used to overcome impotence and raise libido and as a general cure for the disease		Narcotic resin, cannabidiol, cannabidiol-carboxylic acid, cannabigerol and cannabichromene, cannabipinol and cannabidivarin, phloroglucinol β -D-glucoside, tetrahydrocannabinol,	Cannabinoids, Phenol, alkaloid, flavonoid, and volatile oils
<i>Chlorophyllum borivilianum</i> (<i>Asparagaceae</i>)	Safed Musli	In a study of the aqueous extract of dried roots of <i>C. borivilianum</i> in rats, there was increase in libido, sexual vigour and sexual arousal at 250 mg/kg. The study supported treatment of premature ejaculation and oligospermia	The chemical structure of stigmaterol is related to that of testosterone and mainly contributes to its aphrodisiac potentials; hecogenin produces anabolic hormone	Isolated compounds include stigmaterol and hecogenin which are responsible for its antioxidant power, anticancer and aphrodisiac activities. Chlorophytoside-1, fatty acids, eicosadienoic	glycosides, saponins, fatty acids, and hydrocarbons
<i>Citrullus lanatus</i> (<i>Cucurbitaceae</i>)	Watermelon	The effect of red watermelon flesh extract on male sexual behaviour has been determined. In the research, the suspension of the flesh extract was administered on doses 100, 500, and 1000 mg/kg to different groups of male rats (n=5) daily for 22 days. The result showed that oral administration of water melon flesh extract caused significant increase in mounting frequency, intromission frequency and ejaculatory latency. Watermelon flesh extract did not produce undesirable side effects on the male rats and thus its short-term use is apparently safe	Citrulline improves blood drive to the genital regions and plays a significant role in the relaxation of blood, a major tool in high sexual performance	Watermelon contains bioactive agents such as citrulline, β -carotene and lycopene which have been used in the management of prostate cancer	Carotenoids
<i>Eurycoma longifolia</i> (<i>Simarouba-ceae</i>)	Tongkat ali, pasak bumi	Standardized extract F2 at 25 mg/kg and its quassinoids improved rat spermatogenesis, improved testosterone steroidogenesis. standardised water extract at 400 mg/day for six weeks on testosterone, epitestosterone ratio showed significant difference between supplementation and placebo. Treatment with <i>E. longifolia</i> extract at 400 mg/day for 5 weeks resulted to increase in free and total testosterone concentration and muscular force in men and women	Improves spermatogenesis by affecting the hypothalamic-pituitary-gonadal axis. Improves testosterone by inhibiting aromatic conversion of testosterone to estrogen and may also involve phosphodiesterase (PGs) inhibition. The extracts of <i>E. longifolia</i> affects male infertility by suppressing α -2HS glycoprotein expression which thereby increases testosterone level and insulin sensitivity	Quassinoids such as eurycomanone, eurycomnol, pasakbumin-B, hydroxyklaineanonones, β -carboline alkaloids, canthin-6-one alkaloids, eurycomalactone, laurycolactone, biphenyl neolignan and steroids, alkaloids such as 5,9-dimethoxycanthin-6-one, 9,10-dimethoxy-3-methylcanthin5,6-dione have been reported	Phenols, quassinoids, alkaloids, volatile oils, and hydrocarbons

<p><i>Ginkgo biloba</i> (<i>Ginkgoaceae</i>)</p>	<p>Ginkgo</p>	<p>According to some researches, extracts of <i>G. biloba</i> may also help in psychological conditions by easing stress, mild depression and anxiety- major causes of poor sexual performance thereby improving the mood for sexual pleasure. <i>G. biloba</i> extract have been used in traditional Chinese medicine to improve blood circulation. <i>G. biloba</i> constituents have a thinning effect on the blood besides helping to improve the muscle tone in the walls of the blood vessels</p>	<p>Improved blood circulation results to an increase in the amount of oxygen in the blood and to all major organs of the body including the heart and brain thereby resulting to an increased arterial inflow to arterial tissues through arteries and veins without obstructing systemic blood pressure. This enhanced supply of blood to sex organs is crucial in maintaining strong erection</p>	<p>GC-MS, HPLC-MS, HPLC-RI analysis of samples have led to the characterization of ginkgolides A, B, C, J, M with cage structures involving a tertiary butyl group and six membered rings including a spironone system, a tetrahydrofuran and three lactones groups. 33 flavonoids have been isolated from the leaves including amento flavone, quercetin, myricetin, sesquojaflavone, Ginkgetin, Isorhamnetin, etc. Ginkgolic acids have also been isolated; the albumen of the seed also contains neurotoxic 4'-O-methylpyridoxine (ginkgotoxin), etc</p>	<p>Steroids, flavonoid, and ginkgosides</p>
<p><i>Hibiscus sabdariffa</i> (<i>Malvaceae</i>)</p>	<p>Roselle</p>	<p>Pharmacology of the testicular effects of sub chronic administration of <i>H. sabdariffa</i> calyx aqueous extract in rats has been determined. Doses of 1.15, 2.30, and 4.60g/kg for 12 weeks showed in significant change in the absolute and relative testicular weights; significant decrease in the epididymal sperm count and induced testicular toxicity</p>	<p>It decreases the viscosity of the blood and stimulates internal peristalsis</p>	<p>Several compounds have been isolated from different parts of <i>H. sabdariffa</i> including β-carotene, vitamin C, riboflavin, thiamine, and nutrients such as protein, carbohydrates and minerals like calcium and iron. <i>H. sabdariffa</i> is composed chiefly of organic acids, anthocyanins, polysaccharides and flavonoids. Spectroscopic analysis off the aqueous extract of <i>H. sabdariffa</i> have yielded citric acids, hydroxycitric acid, hibiscus acid, malic acid and tartaric acids; oxalic acid as minor compounds. Delphinidin and cyanidin based anthocyanins including delphinidin-3-saambubioside (Hibiscin), cyanidin- 3,5-diglucoside, delphinidin, etc. have been reported</p>	<p>Carotenoids, vitamins, flavonoids, minerals, and amino acids</p>
<p><i>Lepidium meyenii</i> (<i>Cruciferae</i>)</p>	<p>Peruvian ginseng, maca</p>	<p>In a research, treatment of rats with maca at high altitudes prevented high altitude spermatogenic disruption. In a separate study, 1500 mg/kg or 3000 mg/kg orally showed no significant effect on serum levels of leutinizing and follicle stimulating hormone (FSH)</p>	<p>Maca improves stamina and endurance, mood, and libido and erectile capabilities due to the presence of arginine which boosts NO which relaxes blood vessels, the same basic effect Viagra produces</p>	<p>Macamides such as benzylglucosinolate, benzylisocyanate, benzyl nitrile, benzyl alcohol, benzylaldehyde, benzylamine, hexanal, linoleic acid, N-benzylhexadecanamide, alkaloids, fatty acids, amino acids</p>	<p>Macamides, alkaloids, amino acids, and fatty acids</p>
<p><i>Mimosa tenuiflora</i> (<i>Fabaceae</i>)</p>	<p>Jurema preta, calumbi</p>	<p>A research into the spermatoc characteristics of <i>M. tenuiflora</i> on ram showed no significant differences ($P>0.05$) for the progressive motility, spermatoc strength and morphology among the sheep with or without <i>M. tenuiflora</i>. The result indicated that <i>M. tenuiflora</i> does not influence negatively on spermatoc characteristics of the sheep</p>		<p>Two alkaloids have been isolated from <i>M tenuiflora</i> and includes 5-hydroxytyptamine and N, N- dimethyltryptamine. <i>M tenuiflora</i> is also composed of yuremanine and two chalcones; kukulkan A (2',4',-dihydroxy-3'-4-dihydroxychalcone), kukulkan B (2',4',4- trihydroxy-3-methoxychalcone). <i>M tenuiflora</i> is also composed of the steroids campesterol-3-O-β-D-glucopyranosyl, stigmasterol-3-O-β-D-glucopyranosyl and β-sitosterol-3-O-β-Dglucopyranosyl. Saponins such as mimonoside A, mimonoside B, mimonoside C have been isolated. Five 2-phenoxchromones ("uncommon" flavonoids), the tenuiflorin A [5,7-dihydroxy-2-(3-hydroxy-4-methoxyphenoxy)-6 methoxychromone], tenuiflorin B [5,7-dihydroxy-2-(4-hydroxy-3-methoxyphenoxy)- 6-methoxychromone] and tenuiflorin C and 6-demethoxy-4'-O-methylcapillarisin were isolated from the leaves of <i>M. tenuiflora</i></p>	<p>Alkaloids, steroids, and flavonoids</p>

<p><i>Mucuna pruriens</i> (Fabaceae)</p>	<p>Velvet beans, Lyon bean</p>	<p>In different texts of Ayurveda, <i>M. pruriens</i> is most commonly used in aphrodisiac formulations. At 70 mg/kg, treatments significantly improved testosterone quality, ameliorated Psychological stress and improved sperm count</p>	<p>Producing a dose dependent increase in FSH and leutenizing hormone which increases the number of eggs released at ovulation by the action of L-DOPA and dopamine</p>	<p>L-DOPA, serotonin, mucunain, arachidic acid, behenic acid, genistein, glutamic acids, betacarboline, β-sitosterol, cysteine, dopamine, lysine, tryptamine, riboflavin</p>	<p>Alkaloids, amino acids, saponins, and vitamins</p>
<p><i>Musa (Musa paradisiaca/sapientum)</i> (Musaceae)</p>	<p>Banana, plantain</p>	<p>Aqueous extract of <i>M. paradisiaca</i> root on testicular function parameters on male rats at 25, 50 and 100 mg/kg enhanced the testosterone dependent normal functioning of the testes. <i>M. sapientum</i> contain bromine, norepinephrine, dopamine and serotonin in the peel and pulp. Norepinephrine and dopamine elevate blood pressure while serotonin stimulates the blood vessels of the intestine</p>	<p>Increase in blood Circulation</p>	<p>Bromine, rubidium, strontium, saponins, norepinephrine, dopamine, serotonin, vitamin B₆, vitamin a, c and D and natural glucose, fructose. Several compounds such as acyl steryl glycoside such a sitoindoside-I, sitoindoside-II, sitoindoside-III, sitoindoside-IV and steryl glycosides such as sitosterol, <i>myo-inositol</i>-β-D-glucoside have been isolated from fruits of <i>M. paradisiaca</i>, A bicyclic diarylheptanoid, <i>rel</i>-(3<i>S</i>, 4<i>aR</i>,10<i>bR</i>)-8-hydroxy-3-(4-hydroxyphenyl)-9-methoxy-4<i>a</i>,5,6,10<i>b</i>-tetrahydro-3<i>H</i>-naphthol[2,1-<i>b</i>] pyran, and 1,2-dihydro-1,2,3trihydroxy-9-(4-hydroxyphenyl) naphthalic anhydride, 1,7-bis(4-hydroxyphenyl) hepta-4(<i>E</i>), 6(<i>E</i>)-dien-3-one have also been isolated, cyclomusalenol, cyclomusalenone</p>	<p>Saponins, alkaloids, vitamins, glycosides, triterpenes, and sterols</p>
<p><i>Myristica fragrans</i> (Myristiaceae)</p>	<p>Nutmeg, mace</p>	<p>50% ethanolic extract showed significant increase in aphrodisiac properties in mice such as increase in mating frequency, libido and potency. It has also been used in Unani medicine for the treatment of sexual disorders</p>	<p>Stimulation of the nervous system by myristicin</p>	<p>A-pinene, camphene, p-cymene, sabinene, βphillandiene, γ-terpinene, limonene, myrcene, linalool, 3-methyl-4-decan-1-ol, fixed oils like mysristic, stearic, palmitic, oleic and olenolic acids, Licarin B and malabaricone C</p>	<p>Essential oils, fixed oils, and unsaturated aliphatic hydrocarbon</p>
<p><i>Ocimum gratissimum</i> (Lamiaceae)</p>	<p>Ocimum, wild basil</p>	<p>Oral administration of extracts of <i>O. gratissimum</i> at 100, 250 and 500 mg/kg to 6 groups of male rats once a day for seven days showed significant increase in mounting frequency, intromission frequency, erection and aggregate penile reflexes</p>		<p><i>O. gratissimum</i> consist of several essential oils such as thymol, eugenol, methyl charvical, gratissimol, pentoses, hexoses, uronic acid, alkaloids, tannins, flavonoids, methyl eugenol, cis-ocimene, trans-ocimene, pinene, camphor, germacrene-D, trans-caryophyllene, farnesene, l-bisabolone, p-cymene, γ-terpene, α-trans sabinene hydrate, 1,8- cineole, linalool, β-salinene, and geraniol</p>	<p>Volatile oils, alkaloids, and tannins</p>
<p><i>Panax ginseng</i> (Araliaceae)</p>	<p>Ginseng berry</p>	<p>Panax extract standardized with ginsenoside Rg3 significantly produced significant and sustains increase in sexual activity of normal male rats. Improvement in all forms of sexual dysfunction including erectile dysfunction and premature ejaculation</p>	<p>Ginsenosides enhances acetylcholine-induced and transmural nerve stimulation-activated relaxation associated with increasing tissue cGMP mediated by the release of NO</p>	<p>Triterpene glycosides called ginsenosides. Alkanes, alkenes, sterols, fatty acids, carbohydrates, flavonoids, organic acids and vitamin</p>	<p>Saponins, hydrocarbons, flavonoids and vitamin</p>

<i>Passiflora incarnata</i> (<i>Passifloraceae</i>)	Passionflower, wild passion vine	The aphrodisiac effect of the methanolic extract of <i>P. incarnata</i> has been determined in mice. The result showed significant aphrodisiac properties in male mice at all doses- 75, 100 and 150 mg/kg with 100 mg/kg having the highest activity		Several compounds such as flavonoids and other phenolics have been isolated from <i>P. incarnata</i> such as apigenin and luteolin, isovitexin, vitexin, isoorientin, orientin and saponarin. Also isolated from <i>P. incarnata</i> includes schaftoside, isoschaftoside, isovitexin-2'-O- β -glucoside and isoorientin-2-O- β -glucoside	Phenolics, alkaloids, and sugars
---	----------------------------------	---	--	--	----------------------------------

Table 2: Medicinal Plants used for the improvement of sexual performance and virility.

Conclusion

Herbals medicinal plants have a possible to treat the assorted varieties of body ailments. The demand of herbal medicine is increasing day by day in developed yet as developing countries as a result of they are safer and well tolerated as compared to those of allopathic drugs. These plants must be subjected to animal and human studies to figure out their effectiveness in whole organism systems. Many plants have tried helpful within the management of sexual disorders throughout history, even herbs and spices are accustomed increased sexual activities in varied components of the world. There's great would like for substances that are accustomed treat sexual dysfunction in humans. The utilization of aphrodisiacs is outstanding in several countries of the world as well as Asian country like India, China, Sri Lanka, and Pakistan.

Acknowledgement

The authors are thankful to the Mr. Jivan Kumar; Production officer of Kusum Healthcare Pvt. Ltd. Chopanki, Bhiwadi (Rajasthan) India, for her support and helpful in cooperation in the review and data collection process.

Conflict of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

Bibliography

1. Jo JK, et al. "Effect of starting penile rehabilitation with sildenafil immediately after robot-assisted laparoscopic radical prostatectomy on erectile function recovery: A prospective randomized trial". *Journal of Urology* 199.6 (2018): 1600-1606.
2. Goethe VE, et al. "Concordance and discordance of sexual identity, sexual experience and current sexual behavior in 45-year old men: Results from the German Male Sex-Study". *The Journal of Sexual Medicine* 6.4 (2018): 282-290.
3. Lane-Cordova AD, et al. "Association between cardiovascular health and endothelial function with future erectile dysfunction: The Multi-Ethnic Study of Atherosclerosis". *American Journal of Hypertension* 30.8 (2017): 815-821.
4. Jisheng W, et al. "Comparison of TCM and western medicine diagnosis and treatment strategies for erectile dysfunction". *Chinese Canadian Journal of Human Sexuality* 26 (2017): 23-26.
5. Rongmin J, et al. "Meta-analysis of clinical effectiveness of the acupuncture treatment for erectile dysfunction". *Chinese Canadian Journal of Human Sexuality* 27 (2018): 82-85.
6. Kaminetsky JC, et al. "Quality of erections by age group in men with erectile dysfunction". *International Journal of Clinical Practice* 71.10 (2017):12976.
7. Wang X and Wang MLI. "Discussing on the action mechanism of two-way adjustment of acupuncture". *Journal of Sichuan Traditional Chinese Medicine* 35 (2017):40-42.
8. Onyeji IC, et al. "Impact of surgeon case volume on reoperation rates after inflatable penile prosthesis surgery". *Journal of Urology* 197.12 (2017): 223-229.
9. Cocci A, et al. "Predictive factors of patients and their partners sexual function improvement after collagenase clostridium histolyticum injection for Peyronie's disease: Results from a multi-center single-arm study". *The Journal of Sexual Medicine* 15.5 (2018): 716-721.
10. Ren H, et al. "Comparative effects of umbilical cord- and menstrual blood-derived MSCs in repairing acute lung injury". *Stem Cells International* (2018).
11. Wang J, et al. "The safety and efficacy of acupuncture for erectile dysfunction: A Network meta-analysis". *Medicine* 98.2 (2019): 1-4.
12. Burnett AL, et al. "Erectile dysfunction: AUA guideline". *Journal of Urology* 200.3 (2018): 633-641.
13. Pozzi E, et al. "Clinical profile of young patients with erectile dysfunction: Preliminary findings of a real-life cross-sectional study". *European Urology Focus* (2018).
14. Kalka D, et al. "564 Do socioeconomic factors influence the pathogenesis of erectile dysfunction through modifiable risk factors?". *The Journal of Sexual Medicine* 15.7 (2018): 335.

15. Deng C., *et al.* "Treatment of human chronic wounds with autologous extracellular matrix/stromal vascular fraction gel: A STROBE compliant study". *Medicine (Baltimore)* 97.32 (2018).
16. Sarwer DB., *et al.* "4-Year changes in sex hormones, sexual functioning and psychosocial status in women who underwent bariatric surgery". *Obesity Surgery* 28.4 (2018):892-899.
17. Aversa A., *et al.* "Tadalafil improves lean mass and endothelial function in nonobese men with mild ED/LUTS: in vivo and in vitro characterization". *Endocrine* 56.3 (2017): 639-648.
18. Amano T., *et al.* "Administration of daily 5 mg tadalafil improves endothelial function in patients with benign prostatic hyperplasia". *Aging Male* 21.1 (2018):77-82.
19. Mulhall JP., *et al.* "Predictors of erectile function normalization in men with erectile dysfunction treated with placebo". *The Journal of Sexual Medicine* 15.6 (2018): 866-872.
20. Lunsen RHW., *et al.* "Maintaining physiological testosterone levels by adding dehydroepiandrosterone to combined oral contraceptives: II. Effects on sexual function". *Contraception* 98.1 (2018): 56-62.
21. Ahlers CJ., *et al.* "How unusual are the contents of paraphilias? Paraphilia-associated sexual arousal patterns in a community-based sample of men". *The Journal of Sexual Medicine* 8.5 (2011): 1362-1370.
22. McPhail IV. "Age of onset in pedohebephilic interests". *Archives of Sexual Behavior* 47.5 (2018): 1313-1317.
23. Tozdan S., *et al.* "Keep faith in yourself! -A pilot study on the relevance of specific self-efficacy for modifying sexual interest in children among men with a risk to sexually abuse children". *Journal of Sex and Marital Therapy* 6 (2018): 1-14.
24. Corona G., *et al.* "EDEUS, a real-life study on the users of phosphodiesterase type 5 inhibitors: Prevalence, perceptions and health careseeking behavior among European men with a focus on 2nd-generation avanafil". *Sexual Medicine* 6.1 (2018): 15-23.
25. Hutchings DC., *et al.* "Phosphodiesterase-5 inhibitors and the heart: Compound cardioprotection?" *Heart* 104.15 (2018): 1244-1250.
26. Osondu CU., *et al.* "The relationship of erectile dysfunction and subclinical cardiovascular disease: A systematic review and meta-analysis". *Vascular Medicine* 23.1 (2018): 9-20.

Volume 3 Issue 8 August 2019

© All rights are reserved by Saurabh Nimesh., *et al.*