



Barriers to Antiretroviral Medication Adherence among Key Populations in Nigeria

Olaolu Ojenike*

Pharmacist, Society for Family Health Nigeria, Nigeria

*Corresponding Author: Olaolu Ojenike, Pharmacist, Society for Family Health Nigeria, Nigeria.

Received: October 22, 2019; Published: October 24, 2019

DOI: 10.31080/ASPS.2019.03.0428

Abstract

Adherence to antiretroviral medication is vital to achieving viral suppression in people initiated on antiretroviral medication. Adherence is clients' behavior of taking drugs consistently and correctly. Key population who are enrolled on antiretroviral who adhere to their medication always achieve viral suppression.

Keywords: Antiretroviral Medication; HIV; Nigeria

Nigeria has the second largest HIV epidemic within the world and has one of the highest new infection rates in Sub-Saharan Africa [1]. Antiretroviral therapy (ART) is the mainstay within the management of people living with HIV (PLHIV). Adherence to antiretroviral medication is vital to achieving viral suppression in people initiated on antiretroviral medication. Adherence is clients' behavior of taking drugs consistently and correctly. According to Machtinger and Bangsberg [2] "Poor adherence will lead to treatment failure, evolution of drug resistance, and subsequent immunological and clinical failure". More than 95% of the doses got to be taken for best response whereas lesser degrees of adherence are more usually related to virological failure [3].

In all countries and settings, KPs are disproportionately laid low with HIV infection. Sadly, reaching KPs with HIV intervention may well be an enormous challenge as a result of stigma, discrimination, violence, and legislation of KPs lifestyles [4]. Key populations (KPs) include Men who have sex with Men (MSM), people who inject medication (PWID), and Sex workers (SW), people in prisons and closed settings, and transgender people [4]. The purpose of this article is to identify various barriers to antiretroviral adherence and ways to removing these barriers to enhance maximum ART treatment outcomes.

Barriers to ART adherence

The challenges being faced by Key populations (KPs) contribute major barriers to ART adherence among KPs who HIV-positive and consequently poor treatment outcomes are. These barriers which can be patient related, Health-worker related, patient-provider related, treatment related, clinical environment related, have been identified to impede KPs adherence to ART medication. Majority of these can be structural and comprises economic, institutional, political and cultural factors, that collectively influence the extent to which persons living with HIV follow their medication regimens [5]. Examples of institutional factors are logistical barriers, overburdened health care facilities, limited access to mental health services and difficulties in ensuring adequate counseling [5]. Examples of political and cultural barriers are controversies in the provision of treatment for AIDS, migration, traditional beliefs about HIV and AIDS, poor health literacy and gender inequalities [5].

Stigma and discrimination

Internal stigma may be a major barrier to ART adherence among key populations living with HIV. The threat of social stigma might stop folks living with HIV from revealing their standing to others and function a barrier to HIV treatment adherence [6]. concern of stigma and discrimination because of social beliefs, government

legislations, religion, conjointly makes folks living HIV and key population inaccessible to quality health care and this also is a barrier to majority of key populations on ART care. Consequently, there'll be increase in incomprehensible clinical appointments and pharmacy refills that may later results in virologic failure and over-all poor clinical outcomes.

Migration of key population

Mobility of key populations attributable to stigma and discrimination can be a significant barrier to adherence to ART medication. Most men who have it off with men (MSM) and transgender, do migrate to a special location once initiation on treatment. Most female sex workers (FSW) are the foremost full of this issue because of their mobile nature for greener pastures. Also people who inject drugs (PWIDS) conjointly moves to hidden place due to government legislation. Consequently, they missed clinical appointment and pharmacy refills resulting in poor adherence and treatment failure.

Inadequate adherence preparation before initiation on ART

This involves adherence strategic plans before initiation of clients on ART. The reediness of the client will enhance adherence to ART medication. During initial adherence counseling, health workers are obligated to identify major barriers to client adherence by asking questions about client employment, residential location, psycho-social factors, nutrition and other barriers to adherence. This will enhance the health care worker to properly counsel and prepare the client for ART initiation. Inadequate preparation of clients before initial may lead to poor adherence.

Health worker related barrier to adherence

Health workers might additionally contribute to client poor adherence through numerous means of contact with the client. Health workers are expected to attend to ART clients in an exceedingly non-judgmental and indiscriminately manner. This will enhance clients trust and good adherence. Inadequate supply of medicines, poor storage facility may lead to shortage of medicine and this could contribute to poor adherence. Inadequate health care employees in an exceedingly explicit health facility will increase clients' waiting time at the facility and this can discourage client from consistent clinic follow of visits and drug refills. Consequently, this could result to poor adherence and treatment failure.

Interventions to barriers of ART adherence

No single adherence intervention or package of interventions is effective for all populations and every one settings. People's wants

and circumstances may additionally modification over time, and programs and care suppliers thus ought to tailor a mix of possible interventions to maximize adherence to ART supported individual barriers and opportunities. "Programme-level interventions for rising adherence to ART include: avoiding imposing due payments at the purpose of care, mistreatment fixed-dose combination regimens for ART and strengthening drug offer management systems to dependably forecast, procure, and deliver ARV medication and stop stock-outs" [7]. Several individual level adherence interventions are indicated for reasons additionally to rising adherence to ART. as an example, nutritional support, peer support, management of depression and substance use disorders and patient education are important elements of routine health and HIV care.

Conclusion

Absolute adherence to antiretroviral drug is a significant issue for achieving optimum viral suppression in individuals initiated on antiretroviral therapy. Government laws, stigma, discrimination, migration, psycho-social factors contribute to poor adherence among key populations. Consistent advocacy needs to be done to reduce the incidence of stigma and discrimination. Efforts to support and maximize adherence need to begin before ART is initiated. Developing associate adherence prepare and education are very important initial steps. Finally, initial patient education need to introduce basic knowledge regarding HIV, the ARV medication themselves, expected adverse effects, preparing for treatment and adherence to ART.

Bibliography

1. Thompson M., *et al.* "Guidelines for Improving Entry Into and Retention in Care and Antiretroviral Adherence for Persons With HIV: Evidence-Based Recommendations From an International Association of Physicians in AIDS Care Panel". *Annals of Internal Medicine* (2012).
2. Machtinger EL and Bangsberg DR. "Adherence to HIV antiretroviral therapy". *HIV Insite Knowledge Base* (2005).
3. Olowookere SA., *et al.* "Prevalence and determinants of nonadherence to highly active antiretroviral therapy among people living with HIV/AIDS in Ibadan, Nigeria". *The Journal of Infection in Developing Countries* 2 (2008): 369-372.
4. Ugbeno R., *et al.* "Assessment of Antiretroviral Therapy Outcomes among Key Populations after 6 Months of Initiation in Nigeria". *Texila International Journal of Public Health* 6 (2018): 164-176.

5. Kagee A., *et al.* "Structural barriers to ART adherence in Southern Africa: challenges and potential ways forward". *Global Public Health* 6 (2011): 83-97.
6. Rintamaki LS., *et al.* "Social stigma concerns and HIV medication adherence". *AIDS Patient Care and STDs* 20 (2006): 359-368.
7. World Health Organization. "Guidance on operation and service delivery: adherence to ART". Geneva, Switzerland: WHO (2013).

Volume 3 Issue 11 November 2019

© All rights are reserved by Olaolu Ojenike.