



## Phytochemical and Antimicrobial Activity of *Solanum torvum* Against Respiratory Tract Pathogens

M Rajathi D Modilal<sup>1\*</sup>, R Sindhu<sup>2</sup>, R Cecily Rosemary Latha<sup>1</sup> and R Anandan<sup>3</sup>

<sup>1</sup>PG and Research Department of Zoology, Holy Cross College, Tiruchirappalli, India

<sup>2</sup>Department of Biotechnology, Karpaga Vinayaga College of Engineering and Technology, Kanchipuram, India

<sup>3</sup>Department of Computer Science and Engineering, School of Engineering, VELS Institute of Science, Technology and Advanced Studies, Chennai, India

\*Corresponding Author: M Rajathi D Modilal, PG and Research Department of Zoology, Holy Cross College, Tiruchirappalli, India.

DOI: 10.31080/ASPS.2020.04.0465

Received: December 12, 2019

Published: December 26, 2019

© All rights are reserved by M Rajathi D Modilal, et al.

### Abstract

Medicinal plants are extensively used to cure various infectious diseases in human beings, hence *S. torvum* was investigated for its activity against the isolated pathogens. The study was designed to screen and characterize the bacteria isolated from the respiratory tract infected patients. Phytochemical analysis of the plant showed the presence of alkaloids, terpenoids, flavonoids, saponins, steroids and phenols. Aqueous, ethanol and diethyl ether extract of *S. torvum* was prepared; four different concentrations of each extract was taken to determine the antibacterial activity against the isolated bacteria. The ethanolic extract of *S. torvum* showed highest antimicrobial activity in comparison to aqueous and diethyl ether extracts. Thus *S. torvum* has antimicrobial activity and can be used clinically to find novel antibacterial compounds for respiratory tract pathogens.

**Keywords:** Respiratory Tract Infections; Antibacterial Activity; Phytochemical Activity; *S. torvum*

### Introduction

Respiratory tract infections (RTIs), which involve the upper or lower respiratory tract, frequently occurs after birth [1]. RTIs, such as sore throat, earaches, laryngitis, common cold, sinusitis, and mastoiditis, are the most frequently-occurred infections of all human diseases and have been frequently documented [2,3]. RTIs are amongst the most wide spread and serious infections, accounting for over 50 million deaths globally each year. Each year approximately seven million people die as direct consequences of acute and chronic respiratory infection. Bronchitis and pneumonia are the most common infection. Respiratory pathogens like *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* are some of the causative agents responsible for bronchitis and pneumonia [4].

Most respiratory tract infections are caused by viral and bacterial pathogens responsible for higher morbidity and mortality [5]. The leading causes of non-communicable disease deaths in 2008 due to respiratory diseases were 3.9% and 4.2 million deaths were reported due to asthma and Chronic Obstructive Pulmonary Disease globally [6]. Today, nearly 88% of the global population turn to plant derived medicines as their first line of defense for maintaining health and combating diseases. Currently, people of Asia especially India are utilizing plants as part of their routine health

management [7]. Medicinal properties of plants are hinged on the presence of bioactive principles such as alkaloids, phenols, tannins, glycosides and essential oils amongst others [8]. The primary benefits of using plant derived medicines are that they are relatively cheaper than synthetic drugs, offering profound therapeutic benefits and more affordable treatments. Many of the plant materials used in traditional medicine are readily available in rural areas and this has made traditional system of medicine relatively cheaper than modern medicine. As a result, many potent drugs have been purified from plants, including emetin, quinine, artemisin and introduced to modern medical practice.

*Solanum torvum* of the family Solanaceae is commonly known as Turkey berry and cultivated in Africa and West Indies. The plant is sedative and diuretic and the leaves are used as a haemostatic. The ripened fruits are used in the preparation of tonic and haemopoietic agents and also for the treatment of pain. It has antioxidant properties. It is intensively used worldwide in the traditional medicine as poison anti-dote and for the treatment of fever, wounds, tooth decay, reproductive problems and arterial hypertension. *S. torvum* also possesses antimicrobial, antiviral, immuno-secretory, antioxidant, analgesic and anti-inflammatory, anti-ulcerogenic activities, cardiovascular, nephroprotective, antidiabetic, angiotensin and serotonin receptor blocking activities [9].

Therefore in the current investigation, bacteria responsible for respiratory tract infections were isolated from sputum samples of RTI patients and later on characterized. *Solanum torvum* leaves were subjected to extraction with solvents like ethanol, petroleum ether and water. The antimicrobial effect of the prepared extracts were evaluated against isolated respiratory tract pathogens.

## Materials and Methods

### Collection and authentication of plant material

The plant material *S. torvum* (Figure 1) was collected from in and around Chennai, Tamilnadu, India. The plant was authenticated by Prof. Dr. D. Aravind, Department of Medicinal Botany, National Institute of Siddha, Chennai, Tami Nadu, India.



*Solanum torvum*

**Figure 1:** Photograph of the plant.

### Preparation of extracts

Fresh leaves of *S. torvum* was washed thoroughly in tap water and with distilled water and airdried in the shade at room temperature for five days. Shade dried leaves were powdered. The plant powders (100 g) were successively extracted using ethanol and petroleum ether in soxhlet apparatus and crude extraction was done with water. The extracts were dried in vacuum desiccator and were stored in a sterile container for further use.

### Collection of samples

The sputum samples were collected from patient's aseptically in well-labelled sterile, wide mouthed glass bottles with screw cap from Karpaga Vinayaga Institute of Medicinal Science and Government Hospital. Samples were then taken to the laboratory immediately for analysis [10]. On the labels were marked the name, age, sex of the patients and the time of sample collection.

### Isolation and identification of pathogens in sputum samples

The collected samples were processed as per the standard procedure. For isolation and characterization of bacterial flora, the samples were inoculated into Blood Agar (BA) media and incubated at 37°C for 24 hours [11]. Characteristic colonies from the

plates were isolated and then sub cultured to obtain pure culture. All the bacteria were isolated and identified using morphological and biochemical tests adopting standard procedures [12]. Stock culture was maintained in both Agar slant and 20% sterile buffered glycerol.

### Phytochemical screening

The plant extracts were screened for the presence of biologically active compounds like glycosides, phenolic, alkaloids, tannins, flavonoids, saponin and steroids under qualitative analysis; the screening was carried for all the extracts of the plant [13].

### Antibacterial activity of various extracts

Muller-Hinton Agar (MHA) plates were seeded with 24 hours old culture of the isolates. The organic fractions were dissolved in dimethyl sulfoxide (DMSO). Wells were bored using sterilized syringe of pore size 8 mm. Various concentrations of the extracts (250µl, 500µl, 750µl and 1000µl) were added into the sterile 8mm diameter well. Incubation was made at 37°C for 24 hrs. Antibacterial activity was assayed by measuring the diameter of the inhibition zone formed around the well using standard (Hi-Media) scale. The experiment was repeated four times and the average values were calculated for antibacterial activity [14].

## Results and Discussion

### Phytochemical activity

The phytochemicals present in *Solanum torvum* were alkaloids, flavonoids, steroids, terpenoids, Saponins and phenols (Table 1). This is in accordance with an earlier investigation which showed the presence of various phytochemicals such as alkaloids, carbohydrates, reducing sugars, flavanoids, gums and mucilage and proteins [15]. Preliminary phytochemical screening of *S. torvum* indicates the presence of These compounds are known to be biologically active because they protect the plants against infection. Plants are important sources of potentially useful structures for the development of new chemotherapeutic agents. Many reports are available regarding anti-viral, anti-bacterial, anti-fungal, anti-helminthic and anti-inflammatory properties of plants [16].

Sl. No	Phytochemical	Ethanollic Extract	Aqueous Extract	Petroleum Ether extract
1	Alkaloids	+	+	+
2	Terpnoides	+	+	+
3	Saponins	+	+	+
4	Tannins	-	-	-
5	Flavonoids	+	+	+
6	Phlobatannins	-	-	-
7	Steroid	+	-	-
8	Phenols	+	+	+

**Table 1:** Phytochemical Constituents of *Solanum torvum*.

Flavonoids have been referred to as nature’s biological response modifiers because of strong experimental evidence of their inherent ability to modify the body’s reaction to allergen, virus and carcinogens. They show anti-allergic, anti-inflammatory, anti- microbial and anti-cancer activity [17,18]. Tannins are known to possess general antimicrobial and antioxidant activities [19]. Recent reports show that tannins may have potential value as cytotoxic and antineoplastic agents [20]. Other compounds like saponins also have anti-fungal properties. Saponins are a mild detergent used in intracellular histochemistry staining to allow antibody access to intracellular proteins [21]. In medicine, it is used in hyper cholestrolaemia, hyperglycemia, antioxidant, anticancer, anti inflammatory and weight loss, etc. It is also known to have anti-fungal properties [22]. Saponins have been implicated as bioactive antibacterial agents of plants. Plant steroids are known to be important for their cardiotoxic activities, possess insecticidal and anti- microbial properties [23,24]. Plant derived natural products such as flavonoids,

terpenoids and steroids etc. have received considerable attention in recent years due to their diverse pharmacological properties including antioxidant and antitumor activity. Phenolic phytochemicals have antioxidative, antidiabetic, anticarcinogenic, antimicrobial, antiallergic, antimutagenic and anti-inflammatory [25].

The bacteria were isolated and identified using morphological and biochemical tests following standard procedures. Out of the 50 Sputum samples, 23 bacterial isolates were recovered and the biochemical tests revealed that, these isolates belong to 5 species (Table 2). The bacteria isolated from the samples were *Klebsiella pneumoniae*, *Streptococcus pyrogens*, *Escherichia coli*, *Staphylococcus aureus* and *Pseudomonas aeruginosa* were isolated from the sputum samples. In an earlier study following bacteria were isolated from the sputum sample of RTI patients *S. aureus*, *E. coli*, *P. aeruginosa*, *S. pneumoniae* and *K. pneumonia* [26]; this is in accordance with the present investigation except *S. pneumoniae*.

Sl. No	Characteristics	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>S. aureus</i>	<i>P. Aeruginosa</i>	<i>S. pyrogens</i>
1	Shape	Rods	Rods	Cocci	Rods	Cocci
2	Gram Test	-	-	+	-	+
3	Coagulase	-	-	+	-	+
4	Catalase	+	+	+	-	+
5	Citrate	-	+	+	+	+
6	Oxidase	-	-	-	+	-
7	Indole	+	-	-	-	-
8	MR	+	-	-	-	-
9	VP	-	+	+	-	+
10	Motility	Motile	Motile	Non- Motile	Motile	Non- Motile

**Table 2:** Biochemical characterization of isolated bacteria from RTI patients.

**Antibacterial activity**

The results of the antibacterial activities( Table 3) showed that the plant extracts exhibit remarkable activity against the test organisms isolated (*E. coli*, *K. pneumonia*, *P. aeruginosa*, *S. aureus*, and *S. pyrogens*) with zone of inhibition ranging from 5 to 25 mm. From the table it is evident that for the four concentrations taken for each extract; the highest concentration 100 mg/ml showed the

maximum zone of inhibition for the five bacterial species. Ethanolic extract of *Solanum torvum* shows maximum zone of inhibition for *S. aureus* followed by the *S. pyrogens*. Aqueous extract of *Solanum torvum* shows maximum zone of inhibition for *S. aureus* followed by *S. pyrogens*. Petroleum ether extract of *Solanum torvum* shows maximum zone of inhibition for *S. aureus* followed by *S. pyrogens*. Among the three extracts, ethanolic extract exhibited maximum antibacterial activity.

Zone of inhibition (mm)														
S. No	Microorganism	Ethanolic Extract				Aqueous Extract				Petroleum ether extract				Streptomycin (10 mg/ml)
		A	B	C	D	A	B	C	D	A	B	C	D	
1	<i>E. coli</i>	14	10	7	5	15	13	11	8	10	9	7	5	17
2	<i>K. pneumonia</i>	16	11	9	7	16	11	9	7	12	9	10	7	18
3	<i>S. aureus</i>	23	19	15	11	21	17	9	8	15	11	9	5	21
4	<i>P. aeruginosa</i>	15	13	9	7	17	14	11	7	12	13	11	7	16
5	<i>S. pyrogens</i>	19	15	11	9	21	17	8	5	13	9	7	5	20

**Table 3:** Antibacterial activity of leaf extracts of *Solanum nigrum*. (Zone of diameter of three replicates).

A, B, C, D indicates 100, 75, 50 and 25 mg/ml concentrations.

## Conclusion

This study shows that *S. torvum* possess antimicrobial activity against bacteria associated with respiratory tract infectious. The plant can be used as a source of oral drug against respiratory tract infections; however, further studies are required to isolate the active principle from the crude extract for proper drug development.

## Acknowledgment

I am thankful to UGC for the grant received.

## Bibliography

- Sethi S., et al. "Antimicrobial activity of medicinal plants on respiratory tract pathogens". *IJBPR* 4.2 (2013): 148-152.
- Roncevic N., et al. "Treatment of acute infections of the lower respiratory tract in children". *Medicinski Pregled* 55 (2002): 299-304.
- Sazawal S and Black RE. "Effect of pneumonia case management on mortality in neonates, infants, and preschool children: a meta-analysis of community-based trials". *Lancet Infection Disease* 3 (2003): 547-556.
- Ponni V., et al. "Screening of bioactive potentials and phytochemical nature of *Solanum trilobatum* extracts". *Journal of basic and applied biology* 3.3-4 (2000): 134-136.
- Qadri SMH. "Incidence of bacterial respiratory pathogens and their susceptibility to common antibacterial agents". *Journal of the National Medical Association* 85 (1993): 388.
- WHO Library cataloguing-in-publication data: Global status report on non-communicable diseases 68 (2010).
- Samy RP, et al. "Preliminary screening of ethnomedicinal plants from India". *European Review for Medical and Pharmaceutical Sciences* 12 (2008): 1-7.
- Karou D., et al. "Antibacterial activity of alkaloids from *Sida acuta*". *African Journal of Biotechnology* 5.2 (2005): 195-200.
- Jaiswal BS. "*Solanum Torvum*: A Review of Its Traditional Uses, Phytochemistry And Pharmacology". *International Journal of Pharma and Bio Sciences* 3.4 (2012): 104 – 111.
- Kolawale AS., et al. "Prevalence of urinary tract infections among patients attending Dalhatu Araf Specialist Hospital, Lafia, Nasarawa State, Nigeria". *International Journal of Medical Sciences* 1.5 (2009): 163-167.
- Cheesborough M. *District Laboratory Practice in Tropical Countries*, Microbiology second edition Cambridge University Press 2 (2006): 158-195.
- Cowan SJ and Steel KJ. *Cowan and Steel manual for identification of medical bacteria*, 2nd edn Cambridge University Press, London (1974): 176-232.
- Harborne JB. "Methods of extraction and isolation". In: *Phytochemical Methods*. Chapman and Hall, London (1998): 60-66.
- Perez C., et al. "Antibiotic assay by agar-well diffusion method". *Acta Biologiae Ethnomedicinae Experimentalist* 15 (1990): 113-115.
- Kayalvizhi J., et al. "Studies on the Physico-Phytochemical Properties and Hepatoprotective Effect Of *Solanum Torvum* Swartz In CCl<sub>4</sub> Induced Experimental Toxicity In Albino Rats". *International Journal of Pharmacy and Pharmaceutical Sciences* 4.15 (2012): 0975-1491.
- Cushnie TPT and Lamb AJ. "Antimicrobial activity of flavonoids". *International Journal of Antimicrobial Agents* 26.5 (2005): 343-356.
- De Sousa RR., et al. "Phosphoprotein levels, MAPK activities and NFkappaB expression are affected by fisetin". *Journal of Enzyme Inhibition and Medicinal Chemistry* 22.4 (2007): 439-444.
- Rievere C., et al. "Polyphenols isolated from antiradical extracts of *Mallotus metcalifianus*". *Phytochemistry* 70 (2009): 86-94.
- Aguinaldo AM., et al. "Phytochemistry". In: Guevara BQ. (ed.) *A guide book to plant screening phytochemical and biological*. Manila: University of Santo Tomas (2005).
- Mohanta TK., et al. "Evaluation of antimicrobial activity and phytochemical screening of oils and nuts of *Semicarpus anacardium*". *Scientific Research and Essays* 2.11 (2011): 486-490.
- De-Lucca A., et al. "Fungal properties of CAY-1, a plant saponin, for emerging fungal pathogens, 45th inter science conference in antimicrobial agents and chemotherapy abstract (2005): 180.
- Mandal P., et al. "Antimicrobial activity of Saponins from *Acacia auriculiformis*". *Fitoterapia* 76.5 (2005): 462-565.
- Manjunatha BK. "Antibacterial activity of *Pterocarpus santalinus*". *Indian Journal of Pharmaceutical Sciences* 68.1 (2006): 115-116.
- Arts IC and Hollman PC. "Polyphenols are disease risk in epidemiological studies". *The American Journal of Clinical Nutrition* 81 (2005): 317-325.

25. Scalbert A., *et al.* "Dietary polyphenols and the prevention of diseases". *Critical Reviews in Food Science and Nutrition* 45 (2005): 287-306.
26. Abubakar EMM. "Antibacterial efficacy of stem bark extracts of *Mangifera indica* against some bacteria associated with respiratory tract infections". *Scientific Research and Essay* 4.10 (2009): 1031-1037.

**Assets from publication with us**

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

**Website:** <https://www.actascientific.com/>

**Submit Article:** <https://www.actascientific.com/submission.php>

**Email us:** [editor@actascientific.com](mailto:editor@actascientific.com)

**Contact us:** +91 9182824667