

Phytochemical and *In-Vitro* Evaluation of Anti-oxidant Activity of *Mansoa alliacea* Leaves

SK Ameenabee¹, A Lakshmana Rao², P Suguna Rani³, T Sandhya⁴,
N Teja^{5*}, G Ashu⁵, V Bhavya Naga Vani⁶, CH Purna Durganjali⁶ and
N Pavani⁷

¹Associate Professor, Department of Pharmacology, V.V. Institute of
Pharmaceutical Sciences, Gudlavalleru, India

²Professor and Principal, Department of Pharmaceutical Analysis, V.V. Institute of
Pharmaceutical Sciences, Gudlavalleru, India

³Department of Pharmacology, Sri Venkateswara University of Pharmaceutical
Sciences, Tirupathi, India

⁴Department of Pharmacology, Institute of Pharmaceutical Technology, Sri
Padmavathi Mahila Viswavidhyalayam, Tirupathi, India

⁵Department of Pharmaceutics, V.V. Institute of Pharmaceutical Sciences,
Gudlavalleru, India

⁶Department of Pharmaceutical Analysis, V.V. Institute of Pharmaceutical
Sciences, Gudlavalleru, India

⁷Department of Pharmacy, V.V. Institute of Pharmaceutical Sciences, Gudlavalleru,
India

*Corresponding Author: N Teja, Department of Pharmaceutics, V.V. Institute of
Pharmaceutical Sciences, Gudlavalleru, India.

Received: August 13, 2020

Published: September 10, 2020

© All rights are reserved by N Teja, et al.

Abstract

Mansoa alliacea Lam. (Family: *Bignoniaceae*) is a native plant from Amazonian basin in South America. Plant derivatives are used as an anti-inflammatory, anti-oxidant, antiseptic and anti-bacterial. The study was aimed to determine the pharmacognostic and phytochemicals present in *Mansoa alliacea*. Micro and Organoleptic characteristics of fresh and dried leaf samples had been examined. Physicochemical chemical variables have been done by using WHO suggested variables, preliminary phytochemical of leaf sample had been performed to identify the presence of alkaloids, flavonoids, tannins and phenols, and quinones using the ethanolic extract of the leaves of *M. alliacea*.

Keywords: *M. alliacea*; Alkaloids; Flavonoids; Tannins

Introduction

According to the World Health Organization [1], about 65% - 80% of the population in developing countries use medicinal plants to treat their health benefits. *Mansoa alliacea* belongs to the *Bignoniaceae* family, which is used extensively by many of the indigenous peoples of Amazonia. It is commonly referred to as

garlic and Ajossacha [2]. So far, phytochemical studies have shown that plants alkaloids, flavonoids, steroids, tannins and phenols are structurally diverse chemicals. Of modern herbal medicine in S, the plant has also become a popular treatment. America where arthritis, rheumatists, body aches and pain and muscle aches, injuries and pain are widely used. Blooms and flowers are made up of anti

- inflammatory, antioxidant [3] plant antioxidants and anti - bacterial steroids, beta sitosterol, stigmasterol, daucosterol and fucosterol. The *Mansoa* (*Bignoniaceae*) family is a source of compounds containing organosulfur [5]. *M. Alliacea* used for reproductive organs, renal distress, dizziness, epilepsy, sickening cell disease, depression, metabolism, skin grievance, leprosy, impetigo, helminthic infections, foot athlete, tumors. In this study, we are making an effort to standardize *M. The alliacea* field was conducted for the study of leaf morphological, anatomical, physicochemical and preliminary research.

Materials and Methods

Identification and collection of plant material

The whole plant of *Mansoa alliacea* was collected from the surrounding area of Machilipatnam, Andhra Pradesh, India. These plants were identified and authenticated in the Department of Botany, Hindu College, Machilipatnam. The leaves are cleaned and shade dried under room temperature for one week. Then leaves are crushed to powder. The powdered sample was stored in an airtight waterproof container protected from direct sunlight and heat until the powdered sample is used for the extraction process.

Preparation of extracts

The powdered material of *Mansoa alliacea* (leaves) were extracted for 24 hrs with laboratory alcohol(ethanol) in soxhlet apparatus [7,8]. The extracts were concentrated to dryness till free from the solvents.

Phytochemical analysis

Phytochemical analysis [9] of the extract was carried out for the presence of saponins, tannins, flavonoids, alkaloids, glycosides, steroids, carbohydrates, proteins and phenols by different methods.

Hydrogen peroxide scavenging activity

The ability of hydrogen peroxide extracts to scavenge was based on the Ruch11 method. In a phosphate buffer (pH 7.4), a solution of hydrogen peroxide (40 mM) has been made. The hydrogen peroxide concentration was determined by spectrophotometer absorption at 230 nm. Hydrogen peroxide (0.6 ml, 40 mM) was blended with extracts. After 10 minutes, the absorbance of hydrogen peroxide was determined against a blank phosphate buffer solution without hydrogen peroxide. The amount of hydrogen peroxide scavenging calculated as follows by extracts and regular compounds.

Scavenging activity (%) = $[(\text{Abscontrol} - \text{Abssample})/(\text{Abscontrol})] \times 100$

Reducing power assay

The reduction strength was calculated by the Oyaizu12 cycle. Substances with a reduction potential react to potassium ferricyanide (Fe^{3+}) as potassium ferrocyanide (Fe^{2+}), which reacts with ferrous chloride to form a ferrous ferrous system with a maximum absorption of 700 nm. Mixed with a phosphate buffer (2.5 ml) and potassium ferricianide (2.5 ml), one liter of the test solution. The mixture was incubated for 20 minutes at 50°C. The mixture has been applied with trichloroacetic acid (2.5 ml), which was then centrifuged at 3000 rpm for 10 minutes. The top layer of fluid (2.5 ml) was mixed to distilled water (2.5 ml) and a freshly prepared solution of ferric chloride (0.5 ml). At 700 nm, the absorbance was measured. The typical use was ascorbic acid (20 $\mu\text{g}/\text{ml}$). By adding a standard or test content, a blank was prepared. Increased reaction mix absorption indicates an increase in power reduction.

Phosphomolybdenum method

The complete anti - oxidant capacity test is a spectroscopic way to quantify anti - oxidant ability by creating phosphomolybdenum complex 13. In conjunction with 1 ml of reagent (0.6M Sulfuric Acid, 28 mM Sodium and 4 mM ammonium molybdate), take 0.1 ml of the sample solution. The tube is sealed and incubated at 95°C for 90 minutes in a boiling water bath. The absorbance of the solution is measured at 695 nm against blank in UV after the sample has cooled to room temperature. Spectrometer: Spectrophotometer.

Results and Discussion

Preliminary phytochemical analysis

Phytochemical test of *M. alliacea* ethanol extract was done. The existence of saponins, flavonoids, alkaloids, steroids, sugars, and phenolic compounds was shown in *alliacea* leaves (Table 1). As the anti oxidant function is induced by phenolic compounds and the flavonoids, the amounts present in the extract are high, indicating good antioxidant activity. The phenolic and flavonoid scavenging potential is largely attributed to the existence of hydroxyl groups. The presence of phenolic compounds14 in the plant has led to its antioxidant activity and thus its effectiveness as a medication. Flavonoids have been shown to have the impact on membrane permeability and membrane bond enzymes, including ATPase and phospholipase A2, inhibited.

Phytoconstituents	Chemical test	Aqueous extract	Alcohol extract	Chloroform extract
Flavonoids	Shinoda Test	+	+	-
	Zn + HCl Test	+	+	-
	Lead acetate Test	+	+	-
Alkaloids	Dragandroff's test	+	+	+
	Wagner Test	-	-	-
	Hager's Test	-	-	-
Tannins and Phenols	FeCl ³ Test	+	+	-
Saponins	Foaming Test	-	-	-
Steroids	Salkowski test	+	-	+
Carbohydrates	Molisch test	+	+	-
Glycoside	Keller-Killani Test	+	+	-
Amino acids	Ninhydrin Test	+	+	-
Proteins	Biuret Test	+	+	-

Table 1: Phytoconstituents of *Mansoa alliacea* leaf powder.

Phosphomolybdenum method

Total anti-oxidant capacity assay is a spectroscopic method for the quantitative determination of anti-oxidant capacity, through the formation of phosphomolybdenum complex [15]. Take 0.1 ml of the sample solution is combined with 1 ml of reagent (0.6M sulfuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate). The tube is capped and incubated in a boiling water bath at 95°C for 90 minutes. After cooling the sample to room temperature, the absorbance of the solution is measured at 695 nm against blank in U.V. spectrophotometer.

Hydrogen peroxide scavenging activity

Hydrogen peroxide scavenging activity of the ethanol extract and standard was given in table 2 and figure 1. *Mansoa alliacea* ethanolic extract caused a strong dose-dependent inhibition of hydrogen peroxide. At a concentration of 10, 25, 50, 75 and 100 µg/ml of the extract the scavenging percentage was 26.26, 44.85, 48.94, 60.50 and 67.54 respectively.

Concentration	Absorbance at 230nm	Percent inhibition
Control	0.0946±0.0004	-
Standard	0.1850±0.0007	96
10	0.1550±0.0002	28.67
25	0.1419±0.0003	43.91
50	0.1420±0.0016	48.99
75	0.1510±0.0007	62.4
100	0.1597±0.0009	69.58

Table 2: Anti-oxidant activity of *M. alliacea* leaves by H₂O₂ scavenging activity.




Figure 1

Reducing power assay

The reducing power of the extract compared to ascorbic acid is shown in table 3. In the reductive ability measurement, Fe³⁺-Fe²⁺ transformation in the presence of extract sample was investigated. Reducing power assay is a convenient and rapid screening method for measuring the anti-oxidant potential. In this investigation, table 3 and figure 2 shows the reductive capabilities of ethanolic extract of *Mansoa alliacea* when compared to the standard ascorbic acid. The reducing power increased significantly with increasing concentration of the extract. At 10, 25, 50, 75 and 100 µg/ml the reducing power was 30.26, 36.54, 48.09, 56.39 and 62.55 respectively.

Concentration	Absorbance at 230nm	Percent inhibition
Control	0.0901 ± 0.0004	-
Standard	0.1791 ± 0.0017	89.21
10	0.1341 ± 0.0014	33.27
25	0.1421 ± 0.0006	37.25
50	0.1329 ± 0.0021	49.12
75	0.1247 ± 0.0011	57.41
100	0.1135 ± 0.0005	64.57

Table 3: By reducing power assay.




Figure 2

Phosphomolebdenum method

Total anti-oxidant capacity assay is a spectroscopic method for the quantitative determination of anti-oxidant capacity, through the formation of a phosphomolybdenum complex. The activity increased significantly with increasing concentration of the extract was given in table 4 and figure 3. At 10, 25, 50, 75 and 100 µg/ml the Percent inhibition was 30.26, 36.54, 48.09, 56.39 and respectively.




Figure 3

Concentration	Absorbance at 230nm	Percent inhibition
Control	0.0814 ± 0.0002	-
Standard	0.1781 ± 0.0017	89.21
10	0.1261 ± 0.0002	33.27
25	0.1640 ± 0.0004	37.25
50	0.1518 ± 0.0013	49.12
75	0.1331 ± 0.0007	57.41
100	0.1460 ± 0.0014	64.57

Table 4: Phosphomolebdenum method.

Phenolic compounds are known as powerful chain-breaking anti-oxidants, important plant constituents because of their scavenging ability due to their hydroxyl groups and contribute directly to antioxidative action. Phenolic compounds are also effective hydrogen donors, which makes them good anti-oxidant. It is suggested that polyphenolic compounds have inhibitory effects on mutagenesis and carcinogenesis in humans when ingested up to 1gm daily with a diet rich in fruits and vegetables [4,6,10-14].

Conclusion

The current study concludes that the ethanol extract of *mansoa alliacea* leaves contains a considerable amount of flavonoids, and from this we conclude that *mansoa alliacea* leaves have a high anti-oxidant activity. More bioactive compounds isolation would however help to determine their safety and effectiveness as a leading anti-oxidant nominee for pharmaceutical purposes.

Acknowledgement

Authors are sincerely thankful to V.V. Institute of Pharmaceutical Sciences, Gudlavalleru, India for providing necessary facilities for carryout the research work. Chalasani Sudhakar Rao is gratefully acknowledged for helping in the collection of the plant material.

Bibliography

1. World Health Organization. WHO technical report series. Geneva, Switzerland: WHO (2011).
2. Devang Pandya, *et al.* "Pharmacognostic and phytochemical evaluation of aerial parts of Bignonia Alliaceae". *International Journal of Pharmacy and Life Sciences (IJPLS)* 3.1 (2012): 1339-1344 1339.
3. Enriquez M, *et al.* "Antioxidant activity and differentiation of essential oils of Guaviduca (*Piper carpunya* L.) and Sacha Ajo (*Mansoa alliacea* L.)". *Italian Journal of Food Science* 31.5 (2008): 19-28.

4. Govas Llamocca Jenny E., *et al.* "Sterols present in the apolar extract of garlic roots of garlic sacha *Mansoa alliacea*". *Revista de la Sociedad Química del Perú* 84.4 (2018): 513-521.
5. Maria das Graças Bichara Zoghbi, *et al.* "The genus *Mansoa* (Bignoniaceae): a source of organosulfur compounds". *Brazilian Journal of Pharmacognosy* 19.3 (2009).
6. Venkataraman Sowmyalakshmi and Srilakshmi Gunda. "Phytochemical constituents and pharmacological activities of *Kigelia africana* and *Mansoa alliacea* - a comprehensive review". *Asian Journal of Pharmaceutical and Clinical Research* 11 (2018): 29-32.
7. Jeng KCG and Hou RCW. "Sesamin and sesamol: Nature's therapeutic lignans". *Current Enzyme Inhibition* 1 (2005): 11-20.
8. Chandra P, *et al.* "Analgesic and anti-inflammatory activity of methanolic extract from *Jatropha curcas* leaves on experimental animals". *Indian Drugs* 50.8 (2013): 32-38.
9. Li H, *et al.* "Antioxidant activities of extracts and fractions from *Lysimachia foenumgraecum* Hance". *Bioresource Technology* 100.2 (2009): 970-974.
10. Harborne JB. *Phytochemical methods*: London Chapman and Hill Ltd. (1973): 49-188.
11. Gupta M, *et al.* "Anti-ulcer activity of ethanol extract of *Terminalia pallid Brandis* in Swiss albino rats". *Journal of Ethnopharmacology* 97 (2005): 405-408.
12. Keser S, *et al.* "Hydrogen peroxide radical scavenging and total anti-oxidant activity of Hawthorn". *Journal of Chemistry* 2.1 (2012): 9-12.
13. Arulpriya P, *et al.* "In-vitro anti-oxidant testing of the extracts of *Samanea saman* (Jacq.) Merr". *Der Chemica Sinica* 1.2 (2010): 73-79.
14. Prieto P, *et al.* "Spectrophotometric quantitation of anti-oxidant capacity through the formation of the phosphomolybdenum complex: specific application to the determination of vitamin E". *Analytical Biochemistry* 269.2 (1999): 73-79.
15. Evans CA, *et al.* "Anti-oxidant properties of phenolic compounds". *Trends in Plant Science* 2 (1997): 152.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667