



## Nature of Wriggling Action of the Filarial Parasite *Setaria digitata*

Jayakumar K<sup>1\*</sup>, Dhanya N V<sup>2</sup>, Sony Peter<sup>3</sup> and Sree Padma Priya S<sup>4</sup>

<sup>1</sup>Professor and Head, Department of Biochemistry, PMS College of Dental Science and Research, Thiruvanthapuram, Kerala, India

<sup>2</sup>Reader, Department of Biochemistry, Sri Sankara Dental College, Thiruvanthapuram, Kerala, India

<sup>3</sup>Senior Lecturer, Department of Biochemistry, PM S College of Dental Science and Research, Thiruvanthapuram, Kerala, India

<sup>4</sup>Reader, Department of Biochemistry, Sree Mookambika Institute of Dental Sciences, Kulasekhram, Tamilnadu, India

\*Corresponding Author: Jayakumar K, Professor and Head, Department of Biochemistry, PMS College of Dental Science and Research, Thiruvanthapuram, Kerala, India.

Received: November 17, 2020

Published: December 10, 2020

© All rights are reserved by Jayakumar K., et al.

### Abstract

*Setaria digitata*, a cattle filarial parasite, is always in constant motion called the wriggling movement. It is known to take up oxygen even in presence of cyanide. One of the partial reduction products of oxygen formed in the parasite is H<sub>2</sub>O<sub>2</sub>, generated by the parasite specific quinol oxidase called alternative oxidase. The wriggling movement as well as the activity of the parasite specific quinol oxidase and formation of H<sub>2</sub>O<sub>2</sub> was found to be SHAM sensitive. The wriggling movement was also found to be inhibited by iodoacetate a specific inhibitor of SH groups, showing the involvement of SH groups movement of the parasite. Under *in vitro* conditions H<sub>2</sub>O<sub>2</sub> was found to increase the wriggling movement of the parasite and the increase was found to be reversed in presence of added catalase. These findings definitely suggest that the wriggling movement of *S. digitata* is associated with the reduction of oxygen. The process appears to be maintained by the hemoglobin, GSH/GSSG system and the SHAM sensitive quinol oxidase activity.

**Keywords:** *Setaria digitata*; Wriggling Action; Hydrogen Peroxide; Alternative Oxidase

### Abbreviations

SHAM: Salicyl Hydroxamic Acid; H<sub>2</sub>O<sub>2</sub>: Hydrogen Peroxide; Ubiquinones; Q<sub>0</sub>, Q<sub>6</sub> and Q<sub>8</sub>.

### Introduction

*Setaria digitata* is located in the peritoneal cavity of the host, where oxygen tensions are low. It is a lymphatic nematode of the cattle *Bos indicus* and is recommended by WHO [1] as one of the model systems for the study of human filariasis. This parasite is devoid of typical cytochromes [2] and possesses the ubiquinone Q<sub>6</sub> and Q<sub>8</sub> respectively showing electron transport [3] and antioxidant properties [4]. It also possesses many other unique features such as rotenone sensitive and insensitive NADH dehydrogenase

[5] mitochondrial lactate oxidation, presence of glyoxylate cycle, absence of glucose-6-phosphatase etc. Presence of activities of antioxidant enzymes such as superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase and nitric oxide synthase [6] and presence of a number of nonenzymatic antioxidants such as glutathione, α-tocopherol, ubiquinol Q<sub>8</sub> and carotenoids [7] are also reported. *S. digitata* shows high and continuous wriggling activity *in vitro*. This requires enormous amount of energy production and should be supplied continuously. The functional machinery and mechanism of energy production for this puzzling activity and its relevance in the physiology of the worm is not well understood. The studies carried out using H<sub>2</sub>O<sub>2</sub> as modulator may give a possible mechanism and components for generating energy in these parasites.

Parasites in general were once considered to be facultative anaerobes utilizing oxygen when available. Though classes as a facultative anaerobes *S. digitata* takes up oxygen even in presence of cyanide [8], shows parasite specific alternative oxidase activity and  $H_2O_2$  is considered to be a product of this parasite specific oxidase [5]. It also possesses functional hemoglobin like system [9]. The formation of  $H_2O_2$  in living systems as a part of metabolic activity of oxygen. In fact, catalase and other peroxidases utilize  $H_2O_2$  in some cellular oxidation processes leading to several important metabolites [10]. Nematodes including filarial parasites are always in constant motion, known as the wriggling movement. Ubiquinone (Qo) and a known antifilarial drug Diethyl carbamylzine (DEC) have also been shown to inhibit this movement [11,12]. Further study on the wriggling movement of *S. digitata* in response to certain aspects of the metabolism of oxygen forms the subject matter of this paper.

**Materials and Methods**

*Setaria digitata* located in the peritoneal cavity of cattle *Bos indicus* were collected in Tyrode medium [13]. From the local abattoir Tyrode medium has a composition of Sodium chloride 0.8% Potassium Chloride 0.02% Calcium Chloride 0.02% Magnesium Chloride 0.01% Sodium Bicarbonate 0.015%, Sodium Dihydrogen Phosphate 0.05% and Glucose 0.5%. The extraneous materials sticking on the surface of the parasite were removed by thorough washing with the medium. The worms were then kept in tyrode medium at 37 °C until use.

Biochemicals used in the study were purchased from Sigma Chemical Company, USA and all other chemicals used were of high purity.

**Measurement of wriggling score**

In order to find out the role of quinone in designing antifilarial drug the experiments were carried out by incubating the live worms in presence of  $H_2O_2$ , Coenzyme Q, SHAM, and Iodoacetate, glucose oxidase and catalase, were dissolved uniformly in ethanol and water respectively, according to their solubility. Four worms were put in 25. 0 ml of tyrode solution in the presence and absence of inhibitors.

The peristaltic type wriggling movement from one side to the other and back on the medium was scored as one and counts were made for one minute on each organism at the periods indicated is recorded.

**Results**


The normal wriggling score of the worm when kept in Tyrode medium was measured and the values are presented in Table 1.

Maximum wriggling movement was obtained in the first hour followed by a gradual decrease in the score during subsequent hours. The effect of  $H_2O_2$ , a product considered generally to be an end product of cyanide sensitive parasitic organisms [12] and found to be produced in this parasite [3,9] and the wriggling score was examined. With increase in concentration of  $H_2O_2$ , the score was found to increase initially and then dropped and this is clear from figure 1.

Time (hr)	Wriggling score/min
1	8.3 ± 0.13
2	8.1 ± 0.09
3	7.4 ± 0.13
4	7.1 ± 0.07
5	6.8 ± 0.10
6	5.1 ± 0.08

**Table 1:** Normal wriggling rate of *S. digitata* in Tyrode medium. Mean ± SE, n = 6.

The effect of added  $H_2O_2$  in the incubation medium on the wriggling action is given in the fig 1.it is clear from the result that  $H_2O_2$  induces the wriggling movement. As increase in the concentration of the  $H_2O_2$  from 0.015 to 0.025 micromoles increased the wriggling score from 8.3/min to about 12/min. However with further increase in the concentration of  $H_2O_2$  the rate of the movement fell rapidly and the worms got paralyzed.



**Figure 1:** Effect of  $H_2O_2$  on Wriggling Score.

Addition of catalase reversed the effect and the result is shown in figure 3. the paralyzed worm did not show reversal of the effect on addition of catalase. However the enhanced rate of wriggling did come down to normal rate following the addition of catalase in a dose dependent manner.




Figure 2: Effect of catalase on H<sub>2</sub>O<sub>2</sub> induced Wriggling Score.

System	% of wriggling score at 30 min	Nature of worm
A. Tyrode without glucose +Qo(0.16mm)	0	(+)
A + catase	0	(+)
B. Tyrode without glucose +SHAM	0	(+)
B + catase	0	(+)
C. Tyrode without glucose + Iodoacetate	0	(+)
C + catalase	0	(+)
D. control (system minus Q <sub>o</sub> , SHAM, Iodoacetate )	100	(-) Normal score

Table 2: *In vitro* effect of Q<sub>o</sub> , SHAM , and Iodoacetate in the wriggling score of *S. digitata* in different systems.

N = 6 (+) Total paraysis of worm within 10 min,

Q<sub>o</sub> 2,3Dimeithoxy 5 -methyl 1,4 benzoquinone

(-) No paralysis.

The induction of wriggling movement observed with H<sub>2</sub>O<sub>2</sub> could be reversed with the addition of catalase and hence H<sub>2</sub>O<sub>2</sub>, or H<sub>2</sub>O<sub>2</sub> generating systems appears to have a modulatory role in the wriggling movement of the parasite. This is further confirmed by *in situ* generation of H<sub>2</sub>O<sub>2</sub> by glucose oxidase system where H<sub>2</sub>O<sub>2</sub> is released according to the reaction.




Figure 2: *In situ* Generation of H<sub>2</sub>O<sub>2</sub> on the Wriggling movement.

Unlike other intermediates of oxygen, the generation of H<sub>2</sub>O<sub>2</sub> cannot be dismissed as a mere undesirable byproduct because of some of these observations made [14]. It was also observed that even at high concentrations, H<sub>2</sub>O<sub>2</sub> does not induce lipid peroxidation in the parasite [6].

In systems where oxygen uptake is cyanide insensitive, the reduction is carried out by a cyanide insensitive alternative oxidase [15,16] in fact the presence of SHAM sensitive alternative oxidase has been reported in many other cyanide insensitive system particularly in plants. But, unlike *Setaria*, such systems have prominent cytochrome oxidase system. In some it is also a major route of oxidation while in others as in *Setaria* the system is absent or present in significant amounts [15] additionally unlike the plant systems studied, the filarial system reported here is in constant motion and aerobic oxidation takes place via the alternative route [3].

In plant system generally this oxidation is a non-phosphorylating step. However, instances of heat generation is known in some cases and there are recent reports indicating that ROS particularly  $H_2O_2$  is used up by some of the systems [17].

In the *Setaria* system on the other hand it is different and the process is intimately connected with muscle movement as it is clear from SHAM sensitivity. The inhibition of alternative oxidase and the decrease wriggling score in presence of SHAM are proportional. It was already reported from this laboratory that SHAM is also inhibits the formation of  $H_2O_2$  and associated generation of ATP [3]. Lower organisms generally depend on glycolysis for their energy needs and the case of *S. digitata* cannot be different. This is clear from the fact that the major steps in Krebs's citric acid cycle leading to the generation of NADH are missing in *Setaria*. It is true that, though NADH generation is considerably low in *Setaria*, NADPH is relatively high because of the presence of malic enzyme and pentose oxidation [18]. High activities of glutathione peroxidase, glutathione-S-transferase and glutathione reductase indicate a predominant role for the GSH/GSSG system in the metabolism of *S. digitata* [7]. This and the capability of the parasite to synthesize sterols [19] demand a high NADPH level and hence, the latter may not be available for conversion to NADH using the trans-hydrogenases present in the parasite [20]. The presence of hemoglobin system in *Setaria digitata* [9] offers promise for a steady supply of oxygen even though the oxygen tension in the parasite environment is relatively low. Studies with diethyl maleate reveal that depleting glutathione can cause the paralysis of the parasite. The effect of SH-inhibitors on the wriggling movement further confirms the role of thiol compounds in the wriggling movement.

## Conclusion

Hence, it is concluded that the wriggling movement of *S. digitata* is associated with aerobic oxidation leading to the generation of  $H_2O_2$ . The process appears to be maintained by the hemoglobin, GSH/GSSG system and the SHAM sensitive quinoloxidase activity. The oxidation of a protein sulphide to a disulphide can lead to contraction and reduction of the oxidized form can restore the original form, and these two actions can result in the wriggling motion shown by the parasite. The parasite specific quinol oxidase

generally called the alternative oxidase is currently under detailed investigation.

## Acknowledgements

We thank Prof. T. Rama Sharma, Indian Institute of Science, Bangalore for providing the glucose oxidase and for keen interest in the work. Also we thankful to Filaria Control Unit of Thiruvanthapuram District, Kerala for their timely help.

## Conflict of Interest

Authors do not have any conflict of interest for the present investigation.

## Bibliography

1. Hawking F. "Diethylcarbamazine. A review of the literature with special reference to its pharmacodynamics, toxicity, and use in the therapy of Onchocerciasis and other filarial infections". WHO/ONCHO (1978): 78.142.
2. Kaleysa Raj R., et al. "Oxidative activities in mitochondrial like particles from *Setaria digitata*, a filarial parasite". *Biochemistry Journal* 256 (1988): 559-609.
3. Santhamma K R and KaleysaRaj R "Quinone mediated electron transport system in the filarial parasite *Setaria digitata*". *Biochemical Biophysical Research Communications* 174.1 (1991): 386-392.
4. R Abilash Kumar, et al. "Presence of alternative oxidase activity in the mitochondrial like particles from the filarial parasite *Setaria digitata*". *Journal of Parasitic Diseases* 24.1 (2000): 27-30.
5. S Mohan., et al. "The adult specific ubiquinone Q8 functions as an antioxidant in the filarial parasite, *Setaria digitata*". *Biochemical and Biophysical Research Communications* 288.4 (2001): 949-953.
6. S Mohan., et al. "Effect of DEC on scavenger enzymes and lipid per oxidation in *Setaria digitata*". *Biomedicine* 20 (2000): 37-42.

7. Mohan S. "Investigations on the antioxidant systems of the filarial parasite *Setaria digitata*". Ph D Thesis submitted to the University of Kerala (2002).
8. Santhamma K R and Kaleysa Raj R. "Quinone mediated ATP production in the Filarial Parasite *Setaria digitata*". *Biochemical Biophysical Research Communications* 167 (1990): 568-574.
9. Abhilash Kumar R and Kaleysa Raj R. "Presence and formation of heme and occurrence of certain heme proteins in the filarial parasite *Setaria digitata*". *Biochemical and Biophysical Research Communications* 253 (1998): 49-52.
10. Sivan VM and Kaleysa Raj R. "Quinone analogue irreversibly paralyzes the filarial parasite *Setaria digitata* invitro". *Biochemical Biophysical Research Communications* 256 (1999): 81-83.
11. Zachariah T J and Kaleysa Raj R. "Effect of Diethylcarbamazine on lipid metabolism of *Setaria digitata*". *Indian Journal of Medical Research* 83 (1986): 41-52.
12. Bryant C and Behm C. In: "Biochemical Adaptation in Parasites", Chapman and Hall, London. (1989): 51-69.
13. Sugunan VS and Kaleysa Raj R. "Origin of Biologically active excretory secretory materials from *Setaria digitata*". *Current Science* 55 (1986): 331-533.
14. Skulachev VP. "Role of uncoupled and non-coupled oxidations in maintenance of safely low levels of oxygen and its one-electron reluctant". *Quarterly Reviews of Biophysics* 29 (1996): 169-202.
15. Christopher, *et al.* "Solubilization of the alternative oxidase of *cuukoo-pint* (Arunmaeulatum) mitochondria". *Biochemical Journal* (1985): 228. 309-318.
16. Antony L Moore., *et al.* "Control of oxidative phosphorylation in plant mitochondria the role of nonphosphorylating pathways". *Biochemical Biophysical Acta* 1059. (1991): 121-140.
17. Denis P Maxwell. "The alternative oxidase lowers mitochondrial reactive oxygen production in plant cells". *Proceedings of National Academy of Sciences* 96.14 (1999): 8271-8276.
18. Rafi M M and Kaleysa Raj R. "Phosphoenolpyruvate-Succinate-glyoxalate Pathway in the filarial parasite *Setaria digitata*". *Journal of Biosciences* 16.3 (1991): 121-126.
19. Ajitha Kumari G., *et al.* "Biosynthesis of Isoperinoid compounds in the cattle filarial parasite *Setariadigitata*". *Biochemical Biophysical Research Communications* 205 (1994): 24-29.
20. Unnikrishnan LS and Kaleysa Raj R. "Transhydrogenase Activities and malate dismutation linked to Fumarate Reductase System in the Filarial Parasite *Setaria digitata*". *International Journal of Parasitology* 25 (1995): 779-784.

#### Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: [www.actascientific.com/](http://www.actascientific.com/)

Submit Article: [www.actascientific.com/submission.php](http://www.actascientific.com/submission.php)

Email us: [editor@actascientific.com](mailto:editor@actascientific.com)

Contact us: +91 9182824667