

Analytical Techniques for the Determination of Rilpivirine – A Review

Yenda Manishankar* and Mukthinuthalapati Mathrusri Annapurna

GITAM Institute of Pharmacy, GITAM (Deemed to be University), Visakhapatnam, India

***Corresponding Author:** Yenda Manishankar, GITAM Institute of Pharmacy, GITAM (Deemed to be University), Visakhapatnam, India.

Received: November 19, 2020

Published: December 10, 2020

© All rights are reserved by **Yenda**

Manishankar and Mukthinuthalapati Mathrusri Annapurna.

Abstract

Rilpivirine is an Anti-viral drug. Rilpivirine acts by inhibiting the replication of HIV – 1 by binding in a non – competitive manner directly to reverse transcriptase enzyme. In the present paper the authors have presented a review of various analytical methods published so far in the literature for the quantification of Rilpivirine in pharmaceutical preparations as well as biological samples.

Keywords: Rilpivirine; HIV; Anti-viral Drug

Introduction

Rilpivirine is chemically known as 4-[[4-[[4-(E)-2-cyanoethenyl]-2, 6-dimethylphenyl] amino]-2-pyrimidinyl] amino] benzonitrile, is a second-generation non-nucleoside reverse transcriptase inhibitor from antiviral drugs [1,2]. Rilpivirine (Figure 1) ($C_{22}H_{18}N_6$) (Mo. wt. 364.417 g/mol) is used to treat infection caused by human immuno deficiency virus (HIV). Rilpivirine acts by inhibiting the replication of HIV – 1 by binding in a non – competitive manner directly to reverse transcriptase. This leads to conformational changes and altered function of reverse transcriptase. Rilpivirine is associated with a low rate of transient serum aminotransferase elevations during therapy and has been implicated in rare cases of clinically apparent acute liver injury [3].




Figure 1: Structure of Rilpivirine.

Rilpivirine is available in India with brand name Edurant (Label claim: 25 mg) (Chandra Bhagat Pharma Pvt. Ltd) as film coated tablets and Rilpivirine is also available in combination with Dolutegravir (Juluca), Emtricitabine/Tenofovir (Odefsey, Complera) etc. Baert et al developed a long-acting injectable formulation with nanoparticles of Rilpivirine (TMC278) for HIV treatment [4]. The

present review article summarises the analytical techniques so far developed such as spectrophotometry [5-10] (Table 1), high performance liquid chromatography [11-15] (Table 2), liquid chromatography-mass spectrometric methods [16-19] (Table 3) for the determination of Rilpivirine and some of the analytical parameters were highlighted.

Reagent	Linearity ($\mu\text{g/ml}$)	λ_{max} (nm)	Comment	Ref
Bromo thymol blue	4 -20	425	Visible region	[5]
Bromo cresol green		415		
Methanol	2 - 8	282	UV region	[6]
Methanol: Water (8:2)	0.5 – 3.5	305	Multivariate technique	[7]
Methanol	6 - 16	306	First derivative method	[8]
0.01N HCl	0.5 – 7.5	280	Zero order, AUC	[9]
		275-285		
		264		
237	First and Second derivative methods			
DMF: Acetonitrile (10:100)	1-16	281.6	UV region	[10]
Acetonitrile (Diluent)				

Table 1: Review of spectrophotometric methods for the determination of Rilpivirine.

Mobile phase (v/v)/Flow rate (ml/min)/ Detection wavelength (nm)	Column	Linearity ($\mu\text{g/ml}$)	Ref
Acetonitrile: Acetate buffer (pH 4.0) (65:35)/1/260	Develosil ODS HG-5 RP C18 (5 μm , 150 x 4.6 mm)	HPLC (0-25)	[11]
Acetonitrile: 25 mM potassium dihydrogen phosphate (50:50)/0.6/290	Gemini RP C18 (5 μm ; 150 x 4.6 mm)	HPLC (0.025-2.0)	[12]
Acetonitrile: Phosphate buffer (pH 3.5) (60:40)/1/282	C8 (5 μm , 250 x 4.6 mm)	HPLC (10-50)	[13]
Acetonitrile: Potassium dihydrogen phosphate buffer (pH 2.8) with ortho phosphoric acid (40:60)/1/282	Develosil ODS HG-5 RP C18 (5 μm ; 150 x 4.6 mm)	HPLC (0-30)	[14]
0.1% Ortho phosphoric acid: Acetonitrile (65:35)/1/205	Develosil ODS HG-5 RP C18 (5 μm ; 150 x 4.6 mm)	HPLC (20-70)	[15]

Table 2: Review of HPLC methods for the determination of Rilpivirine.

Mobile phase (v/v)/Flow rate (ml/min)	Column	Linearity (ng/ml)	Ref
0.1 mM EDTA in acetic acid: Acetonitrile: Methanol (Gradient mode) 6,7-dimethyl-2,3-dipyridyl) quinoxaline (Internal standard)	Sunfire C18	LC-MS Human plasma (18-715)	[16]
Acetonitrile: Methanol: 0.1% Acetic acid in 5 mM Ammonium acetate (20:25:55)/0.6 Didanosine (Internal standard)	Discovery C18 (5 μm ; 150 x 4.6 mm)	LC-MS/MS sprague dawley rat serum (2-1000)	[17]
Rilpivirine-d6 (Internal standard)	Gemini C18 (150 x 4.6 mm, 5 μm)	LC-MS/MS Human plasma (5-200)	[18]
Acetonitrile and 0.1% formic acid buffer (80:20)/0.5	C18	LC-MS Human plasma (51-200)	[19]

Table 3: Review of LC-MS methods for the determination of Rilpivirine.

Conclusions

The present review article helps the readers to do research in a new field apart from the presenting existing analytical techniques for the anti-viral drug Rilpivirine.

Bibliography

- Garvey L and Winston A. "Rilpivirine: a novel non-nucleoside reverse transcriptase inhibitor". *Expert Opinion on Investigational Drugs* 18.7 (2009): 1035-1041.
- Fernandez-Montero JV, et al. "Rilpivirine: a next-generation non-nucleoside analogue for the treatment of HIV infection". *Expert Opinion on Pharmacotherapy* 13.7 (2012): 1007-1014.
- Weiss J and Haefeli WE. "Potential of the novel antiretroviral drug Rilpivirine to modulate the expression and function of drug transporters and drug-metabolising enzymes in vitro". *International Journal of Antimicrobe Agents* 41.5 (2013): 484-487.
- Baert L, et al. "Development of a long-acting injectable formulation with nanoparticles of Rilpivirine (TMC278) for HIV treatment". *European Journal of Pharmaceutics and Biopharmaceutics* 72 (2009): 502-508.
- Ramya B, et al. "Extractive spectrophotometric method for determination of non-nucleoside reverse transcriptase inhibitor in bulk and pharmaceutical dosage form". *IOSR Journal of Applied Chemistry* 10.11 (2017): 32-40.
- Somsubhra Ghosh, et al. "Method development and validation of Rilpivirine in bulk and pharmaceutical tablet dosage form by using UV – Visible spectrophotometric method". *Asian Journal of Research Chemistry* 5.12 (2012): 1472-1475.
- Vijaya Sri K, et al. "UV-visible spectrophotometric method for the estimation of Rilpivirine hydrochloride in pharmaceutical dosage form by using multivariate technique". *Chemical Science Transactions* 4.3 (2015): 799-805.

8. Vijayalakshmi R., *et al.* "Application of first derivative spectrophotometric method for the determination of Rilpivirine in pure and tablet formulations". *Der Pharma Chemica* 6.1 (2014): 404-406.
9. MasthanammaSkandAlekhyaG. "Development and validation of UV spectrophotometric methods for estimation of rilpivirine in bulk and pharmaceutical formulation". *International Journal of Pharmaceutical Sciences and Research*. 5.2 (2014): 483-489.
10. Girija BB., *et al.* "Development and validation of UV spectrophotometric method for estimation of Rilpivirine hydrochloride in bulk and pharmaceutical formulations". *American Journal of Pharmtech Research* 3.1 (2013): 450-458.
11. Yasodha A., *et al.* "RP-HPLC method development and validation of Rilpivirine". *International Journal of Pharmacy and Analytical Research* 6.1 (2017): 18-38.
12. Abhijit AD., *et al.* "Development and validation of a simple and isocratic reversed-phase HPLC method for the determination of Rilpivirine from tablets, nanoparticles and HeLa cell lysates". *Biomed Chromatography* 29.5 (2015): 709-715.
13. Somsubhra Ghosh., *et al.* "Method development and validation of Rilpivirine in bulk and tablet doses form by RP-HPLC method". *Research Journal of Pharmacy and Technology* 6.3 (2013): 240-243.
14. Raj Kumar B and Subrahmanyam KV. "A Validated Stability-Indicating RP-HPLC method for the determination of Rilpivirine". *Journal of Global Trends in Pharmaceutical Sciences* 5.3 (2014): 1822-1826.
15. Rubesh Kumar., *et al.* "RP-HPLC method development and validation of Rilpivirine pharmaceutical dosage form". *International Journal of Pharmacy and Analytical Research* 8.3 (2019): 329-337.
16. Shibata M., *et al.* "Development and application of a simple LC-MS method for the determination of plasma Rilpivirine (TMC-278) concentrations". *Journal of Medical Investigation* 60.1-2 (2013): 35-40.
17. Raju AV and Appala Raju N. "Development and validation of a LC-MS/MS method for the determination of Rilpivirine in sprague dawley rat serum and its application to pharmacokinetic study". *Asian Journal of Biomedical and Pharmaceutical Sciences* 3.21 (2013): 23-29.
18. Gupta A., *et al.* "Reliable LC-MS/MS assay for the estimation of Rilpivirine in human plasma: Application to a bioequivalence study and incurred sample reanalysis". *Drug Testing and Analysis* 7.4 (2015): 290-299.
19. Laxminarayana., *et al.* "Liquid chromatography-tandem mass spectrometric assay for the non-nucleoside reverse transcriptase inhibitor Rilpivirine in human plasma". *Biomed Chromatography* 27.2 (2013): 172-178.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667