

Influence of Sepiolite's Incorporation in Diet on the Performance and Welfare of Turkey Reared on Two Types of Litter

Naceur M'hamdi^{1*}, Manel Ben Larbi², Aziza Raach Moujahed¹, Ameni Askri¹, Cyrine Darej¹, Khouloud Boukhali¹, Nadia ben Braiek³, Hajer M'hamdi⁴, Kaouther Mrabet³, Abdallah Jebeli⁵

¹National Agronomic Institute of Tunisia, University of Carthage, Tunisia

²Higher School of Agriculture of Mateur, Bizerte, Tunisia

³Ministry of Agriculture. CRDA Sousse, Tunisia

⁴Ministry of Agriculture. CRDA Ben Arous, Tunisia

⁵Société de nutrition animale (SNA), Tunisia

*Corresponding Author: Naceur M'hamdi, National Agronomic Institute of Tunisia, University of Carthage, Tunisia.

Received: May 17, 2019; Published: July 29, 2019

Abstract

This study was carried out to determine the influence of adding sepiolite to the diet on performances of turkey. A total of 960 male broilers of the Hybrid strain were used. The animals received a DFIE starter food crumbled from day 1 to day 28, a growth food1 DF2G granulated from day 29 to day 42, a growth food 2 DF2-3G granulated from day 43 to day 63, a growth food. 3 DF3G pellet day 64 to day 84, and a DF4G finish growth food from day 85 to slaughter (day 119). The chicks have a starting average weight of about 56.8 g. Over all the rearing periods, the standard feed gave the best results in terms of average weight (56.9g) versus (13334.33g), respectively, during periods 1 and 10. The T1 food improved the consumption index and the T2 showed no significant effect. On the other hand, there were no significant differences between the three ST, T1 and T2 treatments and between the two litter types C and PB for the mortality rate. Although no significant difference between the two types of litter (C, PB), the wood chips have the best results in terms of improvement of performance and reduction of poultry diseases.

Keywords: Turkey Flesh; Hybrid Strain; Sepiolite; Zootechnical Performance; Litter

Introduction

Antimicrobial compounds are commonly included in poultry diets for promoting of growth and control of diseases. Feeds containing no chemical additives are increasingly used in poultry nutrition. Therefore, antibiotic growth promoters were discredited by consumers as well as by scientists (Humphrey, et al. 2002). Many scientists searched for alternatives to antibiotic growth promoters [1,2]. Traditionally, clays have been incorporated in diets (10 ± 20 g/kg) as a technological additive (lubricant) to improve feed manufacture [3]. Sepiolite is a feed additive (E-562) used as a binder and anti-caking agent up to 2% in all feeds for all animal species [4]. Sepiolite has effects on the productive performance of broilers and laying hens [5,6] by affecting on the physical and kinetic properties of digesta. Sepiolite may replace growth factors, antibiotics and anticoccidials in poultry diets as therapeutic agents [6-8]. Eser, et al. [9] reported that 1% sepiolite in broiler diets increased body weight gain and reduced the relative weight of abdominal fat and the levels of serum cholesterol and triglyceride. Fernandez, et al. [10] stated the effects of sepiolite on bone

characteristics in broiler. The main objective of this study was to evaluate the effects sepiolites' incorporation on the performance and welfare of turkey reared on two types of litter.

Materials and Methods

Feed Formulation, broilers and experimental design

The trial protocol was conducted during 119 days from 29 December 2015 to 26 April 2016, in the animal nutrition company (SNA) in its experimental station "Ben Khelifa" spread over a perimeter of 3 ha. A total of 960 one-day-old turkeys from the SABA company hatchery were used in this study. Poultry received diets with high-energy concentrations in the order of 2850, 2920, 3025, 3100 and 3200 kcal of metabolizable energy for foods, respectively, starter, growth, fattening and finishing. The three tested diets are ST: standard diet (control), T1: diet with sepiolite in top (1% of EXAL sepiolite) and T2: diet with sepiolite in reformulation obtained by the replacement of 1% of the quantity but introduced in the standard formula by 1% of EXAL sepiolite. Groups of 40 chicks were divided into separate lots randomly selected for one of the experimental treatments. Throughout the trial, commercial manage-

ment procedures were employed, and ambient temperature was controlled. Natural and artificial light was provided on a basis of 23 h for the first 2 days, 16 h from day 3 to day 14, 21 h from day 15 to the slaughter days. All birds were vaccinated against Newcastle disease, Infectious Bronchitis and Gumboro during the second week of their life.

Measured parameters and statistical analysis

The measured parameters are: live weight (LW) at different dates (LW28, LW42, LW56, LW70, LW84, LW98, LW112, LW119), average weights (AW): (AW1-14, AW14-28; AW28-42, AW42-56, AW56-70, AW70-84, AW84-98, AW98-112, AW112-119, average daily gain (ADG): (ADG1-14, ADG14-28, ADG28-42, ADG42-56, ADG56-70, ADG70-84, ADG84-98, ADG98-112, ADG112-119 and ADG1-119), the amount of food ingested (IQ), the consumption index (CI), the mortality rate (TM), the relative humidity of litter (HR) and the health status of turkeys.

The data was analyzed as a 2 × 3 factorial randomized complete block design to determine main effects and interactions, three feed forms (ST, T1 and T2) and two litter types (C, PB). All data were analyzed using PROC GLM of the SAS program [11] (Version 9.1, SAS Institute Inc., Cary, NC). Differences were considered significant at P < 0.05 and the differences between means were separated by least significant difference.

Results and Discussions

Average weight

The average live weight of turkey broilers is similar for all diets. Indeed, the initial weight of 56.8 g to reach a live weight of 13334.33 g at period 10 (table 1). Live weight at day 0 of chicks belonging to the three treatments was statistically identical (P > 0.05). This proves that the lots were homogenous and did not exhibit weight variability.

Period	Average weight (g)	Standard error	Pr > t
P1	56.9229	89.8830	0.5273
P10	13334.3333	89.8830	<.0001
P2	318.9394	92.3460	0.0007
P3	979.3542	89.8830	<.0001
P4	1986.7083	89.8830	<.0001
P5	3640.5833	89.8830	<.0001
P6	5577.7083	89.8830	<.0001
P7	7526.8958	89.8830	<.0001
P8	10076.0000	89.8830	<.0001
P9	12356.0000	89.8830	<.0001

Table 1: variation of average weight during trial.

However, analysis of the variance showed a significant effect (p <0.001) of the diet on live weight (Figure 1). Several authors admit the same results. Castaing [12] showed that the incorporation of

1% sepiolite into the flesh turkey feed leads to an improvement in weight: 5.7% at 6 weeks, 3.9% at 9 weeks and 1.2% at the age of slaughter. Tolsa Group [13] has shown that the addition of a higher level (2%) of EXAL leads to an improvement of 3.23% in total average weight (11.701 kg vs 12.092 kg) against the control diet and 2.69% against other clays.




Figure 1: Effect of feed type on average weight.

In fact, the subjects who received the standard control diet (ST) had significantly greater weights than the groups receiving the treatments (T1) and (T2) (P <0.05) with respective average weights, 5685.45; 5569.51 and 5501.06 g. On the other hand, during the first period (14 days) we recorded average weights of 56.91 and 57.1 g for treatments T1 and T2, respectively, versus 56.92 g for (ST). Whereas during the last week of rearing, the animals fed the regimes with sepiolite in Top, (T1) and reformulation (T2) respectively have average weights of 13334.33 and 9900.42 g against 14803.43g for the batch receiving the standard food control. This was demonstrated by statistical analysis which revealed a significant difference (P = 0.0298) between these three treatments during the study period. This may be due to an interaction between the energy concentration and the presence of EXAL. With the incorporation of 1% of EXAL, growth tends to be 1.03% lower with the least energy-dense food 3100 kcal (T1) and 1.64% with the more energy-dense food. The synergy between the diets and the period has no significant effect on the evolution of turkey average weight. However, we found no effect of litter types (C, PB) on weight (P = 0.3913).

Average daily gain (ADG)

Results shows the evolution of ADG, depending only on the period effect. It is noted that this parameter changes significantly from 60.34% (18.71 vs 47.19 g/d) in start-up period (weeks 1, 2, 3, 4), to 47.99% (138.36 vs 71.95 g/d) during the growing season, 23.53% (182.07 vs. 139.22 g/day) during the fattening period then decreases by 16.52% (162.85 vs 139.76 vs 3.29 g/d) during the finishing period. This coincides with the results obtained by Tolsa Group [13], which has shown a significant increase in turkey weight

gain during all breeding periods. The analysis of the variance revealed no significant difference between the three diets (ST, T1 and T2), and despite the absence of significant effect on the evolution of weight gain ($P > 0.05$) during the whole control period (ST) seems to have a slightly higher weight gain compared to treatments with sepiolite (T1, T2) (Table 2).

ADG (g)	
Feed	
T1	111,99 ± 0,027 ^a
T2	111.46 ± 0,022 ^a
ST	112.49 ± 0,023 ^a
Litter	
PB	111.515 ± 0,21 ^a
C	112.769 ± 0,25 ^a

Table 2: Effects of diet and litter on average daily gain.

In addition, litter types (C, PB) have no significant effect on the evolution of ADG during the four rearing phases ($P = 0.21$) (Table 2). In the same context, Ritez [14], in studying the impacts of straw-berries and straw on the zootechnical performance of turkeys, showed that chopped straws and wood chips have no significant effect on weight gain. Analysis of variance showed the absence of any significant effect of the food-litter interaction (Table 3). But it appears that the ADG recorded for the control (ST) are slightly improved on straw litter in corks compared to the cuttings (115.35 vs. 109.22 g/d) whereas for the treatments with sepiolite (T1, T2), ADG are better for litter animals respectively, 115.35 (T1, C) vs 113.5 g/d (T1, PB) and 113.72 (T2, C) vs 111.28 g/d (T2, PB).

Litter	Diet	ADG (g)
C	ST	109.22 ± 0,364
C	T1	115.35 ± 0,370
C	T2	113.72 ± 0,367
PB	ST	115.75 ± 0,374
PB	T1	

Table 3: Effect of the interaction feed-litter on average daily gain.

Ingested feed (IF)

The amount of food ingested varies significantly with changes in turkey age ($P < 0.0001$). Indeed, we recorded significant improvements towards the third period of 1.18% (22.95 vs 35.91 g), the sixth period of 5.94% (69.5 vs 134.26) g and towards the ninth period of 5.83% (167.33 vs 230.94 g). The daily consumption of animals is in line with the application of the energy ration plan. The difference in energy concentration of 200 kcal between foods (ST and T2) leads to a significant difference ($P < 0.001$) of average consumption greater than 1.15%, (124.83 g/floor/period vs 120.65 g/floor/period). Foods with sepiolite in top (T1) allowed to consume a quantity lower than the control of 1.9% (124.83 g/office/period vs 117.74 g/office/period). Knowing that the same daily energy

intake was assured with all foods. These results confirm those obtained previously under similar conditions [12,15]. The litter had no effect on the quantity of food ingested, but it showed, in synergy with the different diets (ST, T1, T2), significant differences ($P < 0.05$). In fact, feed consumption (ST and T2) by turkeys reared on straw corks improved by about 0.7% and 2.42%, respectively. Only the regimen with sepiolite in top (T1) shows an improvement on wood chips of 2.16%. In the same context, many studies admitted the lack of significant litter effect on performance while others showed that the wood chips [12,15]. In contrast, others studies showed the significant effect of both litter types (C, PB) on flesh poultry performance [13,16].




Figure 2: Variation of ingested food with diet types.

Consumption indice

For the evolution of food consumption per week, we observe a significant increase ($P < 0.001$) in the consumption index over all rearing periods. These fluctuations in C-teeth can be explained by the presence of a latency time necessary for the turkey to adapt after each food transition (starter food. growth food. fattening feed and finishing food).

The type of food has no significant effect on the consumption index ($P = 0.22$). Besides, there is no significant difference between the three diets (ST, T1, T2). The incorporation of 1% sepiolite in top or reformulation into the fleshed turkey feed has no significant effect ($P > 0.005$) on palatability (Table 4). On the other hand, there is a slight non-significant increase ($P = 0.22$) in the consumption index for the lots receiving the control diet (ST). (2.47) against the batches receiving the T1 and T2 regimens which respectively (2.377 and 2.432).

Our results disagree with those of Castaing, *et al.* [17]. The incorporation of 2% Sepiolite in turkey food significantly improves the consumption index with a rate of 4.3%. This can be explained by the fact that the addition of sepiolite at a rate of 1% in the diet can never pass without leaving effects on the feeding behavior of the animals. In fact, the food that contains sepiolite is often characterized by a slightly increased hardness compared to the control food and its ability to slow food transit. Besides the bibliographies provide the best feed efficiency is guaranteed for the most energy

intake which coincides in our case. The diet with sepiolite in top (ST) is in cumulates on all breeding periods more consumed. The litter type did not show any significant effect (P = 0.082) on the turkey consumption index.

Diet	Consumption indice
ST	2.470 ± 0.226 ^a
T1	2.377 ± 0.228 ^a
T2	2.432 ± 0.226 ^a

Table 4: Effect of feed type on consumption indice.

Mortality rate

Mortalities are recorded every day at each level during the breeding period and at the end of the experiment collected per week. During all the breeding periods, mortality rate varies significantly (p <0.001). Indeed, during the first phase of growth that the mortality rate is highest during the 17 weeks exceeding 6, 04% (Figure 6). Mortality was recorded in the first few days after litter renewal. especially in straw and cork flooring. The analysis of cadavers showed that this increase in mortality due mainly due to litter consumption. The diet has no significant impact on the mortality rate (P> 0.05) (Table 5). The litter has not a significant impact on the mortality rate (P = 0.34). This is confirmed by previous studies that showed that litter has no direct effect on turkey mortality rate although it may be origin of certain pathologies [18,19].

Mortality (%)	
Diet	
ST	1.11 ± 0.420 ^a
T1	1.92 ± 0.424 ^a
T2	1.49 ± 0.424 ^a
Litter	
PB	1.50 ± 0.34 ^a
C	1.51 ± 0.34 ^a

Table 5: Effect of diet and litter types on mortality rate of birds.

Health status of birds

During breeding period (19 weeks), the bulbs at the keel bone level changes significantly (P <0.001) with the turkey age (5%), this can be explained by the accumulation of waste and droppings and the lowering of building hygiene status and the higher of litter moisture level. Birds reared on wood chunks had significantly lower (P <0.001) keel bone infections and ampoules (0.53%) than those raised on straws. It is therefore the cutters that seem to allow the best results in terms of slowing down the evolution of the keel bulbs. These results confirm the positive effects of the use of the cuttings on the performances of birds [14], notably by decreasing litter crusting [16].

Conclusion

This study aims to use food additives and litter in poultry farming. Our results showed the effectiveness of additives and bedding as a promising approach to poultry farming to improve feed quality, growth performance and bird health status. In this study, we used sepiolite as a technological additive and straw straw as litter. To this end, three diets were studied on a large turkey population on two types of bedding to determine their effectiveness from the first day to age at slaughter. The ST standard control diet allowed the highest zootechnical performance in terms of average weight and average weight gain. The diet with sepiolite in top T1 allowed the best consumption index and the ration with sepiolite in reformulation has no improvement effect (P>0.01). The two regimens with sepiolite did not show any effects on the mortality rate for the different litter types. For the litter, the cuttings have the best results in terms of slowing of the evolution of the cornet bulbs. While, it does not show any significantly effect on the buttons of the breast, the scratches and the lesions of foot pad. Straw plugs have no effect on zootechnical performance and on health status during the four rearing phases.

Bibliography

- Langhout P. “New additives for broiler chickens”. World Poultry-Elsevier 16.3 (2000): 22-27.
- Kamel C. “Tracing modes of action and the roles of plant extracts in non-ruminants”. in: Recent advances in animal nutrition. P.C. Garnsworthy and J. Wiseman, Eds. Nottingham University Press, Nottingham (2001): 135-150.
- Angulo E., et al. “Effect of sepiolite on pellet durability in feeds differing in fat and fibre content”. *Animal Feed Science and Technology* 53 (1995): 233-241.
- EFSA. European Food safety Authority (2015).
- Ouhida I., et al. “Enzymes (β-glucanase and arabinoxylanase) and/or sepiolite supplementation and the nutritive value of maize-barley wheat-based diets for broiler chickens”. *British Poultry Science* 41 (2000): 617-624.
- Tortuero F., et al. “Effects of dietary sepiolite on the growth, visceral measurements and food passage in chickens”. *Archives of Zootechnology* 41 (1992): 209-217.
- Ayed MH., et al. “Effect of sepiolite supplementation on broiler growth performances and carcass yield. Proc. West. Sect”. *American Society of Animal Science* 59 (2008): 169-172.
- Viseras C and Lopez-Galindo A. “Pharmaceutical application of some Spanish clays (sepiolite, palygorskite, bentonite): some pre formulation studies”. *Applied Clay Science* 14 (1999): 69-82.

9. Eser H., *et al.* "Effects of sepiolite usage in broiler diets on performance, carcass traits and some blood parameters". *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 18 (2012): 313-318.
10. Fernandez E., *et al.* "The effects of different levels of dietary sepiolite on tibial dyschondroplasia in chickens". *Archives of Geflügelk.* 58 (1994): 171-175.
11. Statistical analysis system. SAS user's guide. Version 9.0. Cary, SAS. Institute, (2002).
12. Castaing J. Effect de l'inclusion de 2% de Sepiolite "EXAL" dans les aliments à deux niveaux énergétiques présentés en granulés pour porcelets et porcs charcutiers". Journées de la Recherche Porcine en France 21 (1989): 51-58.
13. Tolsa group. EXAL T: Optimizing binder granulation additive feed and the yield of the plant (2002).
14. Ritz CW., *et al.* "Implications of ammonia production and emissions from commercial poultry facilities: A review". *Journal of Applied Poultry Research* 13 (2004): 684-692.
15. Parisini SL., *et al.* "Effetti della sostituzione con sepiolite di parte de la mangime nelfalimentazione del suino pesante". Atti10 et dquo, Congresso Nazionale ASPA (1993): 459-464
16. Tasistro AS., *et al.* "Spatial variability of broiler litter composition in a chicken house". *Journal of applied poultry research* 13 (2004): 29-43.
17. Castaing J., *et al.* "Incidence de deux niveaux d'apports en vitamines sur les performances du poulet de chair". Cinquièmes Journées de la Recherche Avicole, Tours (2003).
18. Castaing J. "Uso de las arcillas en alimentación animal". In: CURSO DE ESPECIALIZACIÓN, 14, 1998, Barcelona. Avances em nutrición y alimentación animal. Barcelona: Fundación Española para el Desarrollo de la Nutrición Animal (1998): 141-158.
19. Folegatti E., *et al.* "Prevalence of foot pad lesions and carcass injuries as indicators of broiler welfare conditions in Italy". In: 12th European Poultry Conference, Verona, Italy (2006): 303-307.

Volume 1 Issue 1 August 2019

© All rights are reserved by Naceur M'hamdi, *et al.*