



Improving Livestock Production Using Auxiliary Reproductive Biotechnology

Talha Safdar¹, Muhammad Ammun Bashir¹, Hamza Jawad^{1*}, Nasir Iqbal², Ayesha Sadiq³, Zubair Luqman¹, Sadaf Aslam², Naveed Hussain², Suwaibatul Annisa⁴ and Hiba Qasir⁵

¹Faculty of Veterinary and Animal Sciences, The Islamia University of Bahawalpur, Bahawalpur, Pakistan

²Department of Veterinary Surgery and Pet Sciences, University of Veterinary and Animal Sciences, Lahore, Punjab, Pakistan

³Department of Pathology, University of Veterinary and Animal Sciences, Lahore, Punjab, Pakistan

⁴Faculty of Veterinary Medicine, Airlangga University, Indonesia

⁵Center for Integrated Breeding Research, Faculty of Agriculture Sciences, Georg-August-Universität Göttingen, Göttingen, Germany

*Corresponding Author: Hamza Jawad, Faculty of Veterinary and Animal Sciences, The Islamia University of Bahawalpur, Bahawalpur, Punjab, Pakistan.

DOI: 10.31080/ASVS.2020.02.0078

Received: June 05, 2020

Published: June 30, 2020

© All rights are reserved by Hamza Jawad, et al.

Abstract

From the beginning, many reproductive techniques have been used in domestic animals to achieve the desired goals. These goals include: Increasing reproductive and productive efficiency, improved genetic makeup of farm animals, reducing the risk of diseases and injury, producing resistance against climate and infectious agents, and research. Major Assisted Reproductive Techniques (ART) discussed in this brief review are, Artificial Insemination (AI), Estrus Synchronization, Superovulation, Semen Sexing, Cryopreservation, and Embryo Transfer (ET). Artificial insemination is the most basic and widely used technique for improving genetics and increasing conception rate. Estrus synchronization is the technique in which the selected females are targeted to come in heat (estrus) simultaneously in a brief period using different protocols. Super-ovulation is producing multiple ova at a single time using many combinations of hormones. Semen Sexing, as the name implies, is a method in which sperms are sorted out, using different techniques, to gain animal of desirable sex. In Cryopreservation, long-term preservation of living cells and tissues is done at a very low temperature (conventionally -196°C) without compromising the structural and functional integrity of the preserved specimen. Embryo transfer involves producing and collecting embryos in donor animals and then implanting them in recipients. Reproduction is the backbone of any livestock enterprise, and to achieve remarkable results, the use of Assisted Reproductive Techniques (ART) is inevitable. This mini review article will provide a brief comparison of such great reproductive biotechnological methods which are using in livestock farming.

Keywords: Artificial Insemination; Estrous Synchronization; Super Ovulation; Semen Sexing; Cryopreservation; Embryo Transfer

Introduction

Assisted breeding techniques are key to improving animal production efficiency in livestock businesses. Unfortunately, these techniques are being used in only a few farms to maintain production levels [1]. These technologies aim to eradicate common reproductive problems in the dairy industry like infertility, low conception rate, less production, the poor genetic makeup of animals, and

early embryonic mortality [2]. The livestock industry is suffering from a huge economic loss because of little use of these beneficial techniques [3]. This article targets to cover an overview of major and easily applicable assisted reproductive techniques (ART) and why there is a need for using all these technologies in today's world [10,11,14].

Artificial insemination

Artificial insemination (AI) is the first and basic assisted reproductive technique that is now being practiced all over the world, including developed and under-developed countries [9,10]. Artificial insemination (AI) is the unnatural depositing of semen in the female reproductive tract and was first practiced in bitch by Spallanzin in the year 1784. Later on, Ivanoff (1922) also used this technique in domestic animals including horses, poultry, cattle, and sheep. Nowadays, almost all corporate and commercial dairy farms are using AI in a routine. Every year almost 100 million cattle, 40 million pigs, 33 million sheep, and 0.5 million goats are artificially inseminated [12]. AI is practiced less in small ruminants, especially goats, as compared with large ruminants (cattle and buffalo). AI involves a collection of semen from the male by any method (most commonly artificial vagina and electro-ejaculation methods), examining its vigor and motility, processing it, and then loading it into the straws. Either freshly chilled or frozen-thawed semen can be used to inseminate the animal [15]. Although Freezing and long-term storage require the use of cryoprotectants. It is also noted that the conception rate is low if the frozen-thawed semen is used. AI is an easy way to improve the genetic makeup of progeny from the side of a male. Also, there is an improved conception rate, less chance of injury, and a low risk of spreading sexually transmitted diseases (STD) [4].

Estrus synchronization

The sole purpose of estrus synchronization is to induce estrus in a desired group of females at a predestined date and time. While synchronizing the time of induction of heat, we are ultimately scheduling the time of parturition. Detection of heat is a crucial part of the good management of animals and this is even harder when we have to deal with a large herd of hundreds and thousands of animals. Using the Estrus synchronization technique, relieve from this difficulty by bypassing the estrus detection phase. The most commonly used approaches for synchronizing the ovulation time are Prostaglandin Treatment, Controlled internal drug-releasing regimen, and GnRH-PGF2 α -GnRH protocols. During Prostaglandin treatment we are using a luteolytic agent (PGF2 α) 11 days apart in cyclic females regardless of the stage of estrus in animals and then after the appearance of estrus signs, timely AI is performed. CIDR (controlled internal drug-releasing) device maintains the blood progesterone concentration in the targeted animals and is inserted into the vagina for 7 days. One injection of PGF2 α is used on day 6 and estrus is seen on day eight. GnRH-PGF2 α -GnRH protocol is a good regimen to induce estrus in non-cyclic animals. Usually on day 1 GnRH is injected, and then 7 days later, PGF2 α injection is ad-

ministered. Another injection of GnRH is given on the 10th day and AI is performed on day 11. The use of estrus synchronization technique eliminates the possibility of human error in the detection of estrus. It is also very beneficial for animals, which exhibit low external heat symptoms (silent heat) like sheep and buffalo [2,3,7].

Superovulation or multiple ovulation

As the name superovulation or multiple ovulations indicates, it is the induction of ovulation of multiple ova from the ovary of an animal using different injectable hormones [8]. Any cyclic animal can be made to ovulate multiple ova at a single time using synthetic follicle-stimulating hormone (a gonadotropin which is naturally released from the anterior lobe of the pituitary gland) [11,17]. Multiple ovulation techniques combined with embryo transfer can bring revolution in the livestock industry. Although superovulation can be induced in any cyclic animal it is usually applied in animals of superior genetic. As Artificial insemination can be used for the improvement of genetic from the male side, multiple ovulations is being used as an improvement of genetics from the female side [18].

Usually, a follicle-stimulating hormone is injected at the end of the luteal phase or before 1-2 day of completion of the estrus synchronization regimen [14,15]. Superovulation is a crucial technique for embryo biotechnology. Several factors should be considered before inducing superovulation in an animal. Most important of these are breeds, reproductive health status of the animal and nutrition. Also, repeated super ovulations have a significant effect on the next one [16,17].

Semen sexing

For producing progeny of desired gender, either male or female, the technique of semen sexing is used. There is a difference in the DNA content of X chromosome bearing sperm and Y-Chromosome bearing spermatozoa [5-18]. Semen sexing is done by flow cytometric separation of fluorescent-labeled X-Chromosome possessing spermatozoa and Y-Chromosome possessing spermatozoa [31]. In recent times, this technology was capable of sorting 15 million spermatozoa into X and Y chromosome bearing chromosomes [5,19]. The accuracy of predicting the sex of the resulting progeny is 85-95 percent. Although new generation flow cytometers with the high efficiency of sorting of sperms are also being devised and utilized nowadays. There are some flaws in the sexing of semen like low sperm concentration, reduced motility, viability, decreased life span of spermatozoa, and some structural defects. Also, sexed semen needs deep intra-uterine insemination [20]. But still, the benefits of this technique outnumber the drawbacks. Major advantages include a high ratio of progeny testing programs, enhanced male

production for this purpose, low risk of sex-limited diseases, and the last but not least, conservation of superior and rare genetics of animals [5,21].

Cryopreservation

It is a technique in which living cells and tissues are preserved in a state of suspended animation at a very low temperature (-196 °C) for a long period while maintaining the structural integrity of the specimen [6,22]. At a very low temperature, the processes going on in a living cell are dramatically reduced [13]. To alter these processes different cryoprotective agents (CPA) are used like Glycerol, Dimethyl sulfoxide (DSMO), and some polymers. Liquid Nitrogen is used as a freezing agent [1,8]. Some difficulties like ice crystal formation at a fast cooling rate and osmotic changes due to the high concentration of intracellular and extracellular solutions at a low cooling rate were faced in conventional methods of freezing [23]. These problems can lead to cryo-injury. But now, vitrification has reduced the risk of cryo-injury very much. Cryopreservation of sperms, oocytes, and embryos has brought a revolution in breeding programs and research systems. Other benefits of cryopreservation are cryosurgery, preservation of organs and tissue for transplantation, and application in food sciences [5,29].

Embryo transfer

Embryo Transfer is a technique which can improve the genetic worth of a herd at a faster rate than any other techniques discussed above. Embryo Transfer is a process that involves many steps. Mainly, it involves generating and obtaining embryos from females of superior genetics called as the donor and then implanting these embryos after processing and grading, into the uterus of different females known as recipients [25,26]. ET is an expensive technique and is only practiced in animals which can justify its cost. While producing embryos in the donor females, estrus synchronization and superovulation methods are applied to gain better results [26]. By using this technique, genetic combinations of both males and females can be used to enhance genetic gain in the offspring. Surgical, as well as non-surgical embryo transfer techniques, are being practiced in elite cattle. Although, non-surgical embryo transfer is cost-effective, safe, and shows better results [19,30]. This technique is not much common as other assisted reproductive techniques. It requires highly skilled technicians and embryologists. The International Embryo Transfer society provides technicians for this purpose. This is the best way devised yet for the rapid expansion of animals of elite genetic makeup [27,28].

Conclusion

The use of auxiliary breeding techniques in farm animals has increased over the past few decades. With time assisted reproduc-

tive techniques are drawing attention and becoming popular in the livestock industry due to astonishing results. Yet many technical refinements and public awareness are needed in this field. Artificial Insemination, superovulation, semen sexing, cryopreservation and embryo transfer have proved to be valuable in improving the genetic merit of animals of low genetic worth. Estrus synchronization has eliminated many managerial problems in the reproduction of farm animals. New technologies like stem cell culture technique, Intra-cytoplasmic sperm injection and nano-technology are also being trialed. Therefore, in order to deal with the future crisis, there is an urgent need to expand reproductive biotechnology with large and basic support at all levels around the world.

Bibliography

1. Verma OP, *et al.* "Assisted Reproductive Techniques in Farm Animal-From Artificial Insemination to Nanobiotechnology". *Veterinary World* 5.5 (2012).
2. Gelayenew Bizelew and Getahun Asebe. "Review on Major Assisted Reproductive Technologies". (2016).
3. Gelayenew B and Asebe G. "Review on Major Assisted Reproductive Technologies" (2016).
4. Agca Yuksel and Hongsheng Men. "Gender Selection in Mammalian Semen and Preimplantation Embryos". *Comparative Reproductive Biology* (2007): 147-157.
5. Choudhary KK, *et al.* "Advances in reproductive biotechnologies". *Veterinary world* 9.4 (2016): 388.
6. Woods Erik J, *et al.* "Fundamental cryobiology of reproductive cells and tissues". *Cryobiology* 48.2 (2004): 146-156.
7. Schatten Heide and Gheorghe M. "Constantinescu, eds. Comparative reproductive biology". Blackwell Publication (2007).
8. Tsai S and C Lin. "Advantages and applications of cryopreservation in fisheries science". *Brazilian Archives of Biology and Technology* 55.3 (2012): 425-434.
9. Chakravarthi P, *et al.* "Use of assisted reproductive technologies for livestock development". *Veterinary World* 3.5 (2010).
10. Durrant BS. "The importance and potential of artificial insemination in CANDES (companion animals, non-domestic, endangered species)". *Theriogenology* 71.1 (2009): 113-122.
11. Mermillod Pascal. "In vitro production of ruminant embryos: results, limits and perspectives" (2007).

12. Boa-Amponsem K and G Minozzi. "The state of development of biotechnologies as they relate to the management of animal genetic resources and their potential application in developing countries". Background Study Paper 33 (2006).
13. Hiemstra, *et al.* "The potential of cryopreservation and reproductive technologies for animal genetic resources conservation strategies". The role of biotechnology in exploring and protecting agricultural genetic resources. FAO, Rome (2006): 45-59.
14. Paterson Lesley, *et al.* "Application of reproductive biotechnology in animals: implications and potentials: applications of reproductive cloning". *Animal Reproduction Science* 79.3-4 (2003): 137-143.
15. Vishwanath R. "Artificial insemination: the state of the art". *Theriogenology* 59.2 (2003): 571-584.
16. Squires EL, *et al.* "Embryo technologies in the horse". *Theriogenology* 59.1 (2003): 151-170.
17. Stenbak TK, *et al.* "Ovulation rate in ewes synchronized with Syncro-Mate-B (SMB) and follicle stimulating hormone". *Small Ruminant Research* 48.1 (2003): 1-8.
18. E Silva J Chagas, *et al.* "Plasma progesterone profiles, ovulation rate, donor embryo yield and recipient embryo survival in native Saloia sheep in the fall and spring breeding seasons". *Theriogenology* 60.3 (2003): 521-532.
19. Hasler John F. "The current status and future of commercial embryo transfer in cattle". *Animal Reproduction Science* 79.3-4 (2003): 245-264.
20. Gonzalez-Bulnes A, *et al.* "Effect of follicular status on superovulatory response in ewes is influenced by presence of corpus luteum at first FSH dose". *Theriogenology* 58.8 (2002): 1607-1614.
21. Stachecki James J, *et al.* "Fetal development of mouse oocytes and zygotes cryopreserved in a nonconventional freezing medium". *Cryobiology* 44.1 (2002): 5-13.
22. Maclellan LJ, *et al.* "Pregnancies from vitrified equine oocytes collected from super-stimulated and non-stimulated mares". *Theriogenology* 58.5 (2002): 911-919.
23. Terawaki Y and Y Asada. "Relationships between distribution of number of transferable embryos and inbreeding coefficient in a MOET dairy cattle population". *Asian-Australasian Journal of Animal Sciences* 15.12 (2002): 1686-1689.
24. Mitchell LM, *et al.* "Season affects characteristics of the pre-ovulatory LH surge and embryo viability in superovulated ewes". *Animal Reproduction Science* 74.3-4 (2002): 163-174.
25. Stenbak TK, *et al.* "Effects of follicle stimulating hormone (FSH) on follicular development, oocyte retrieval, and in vitro fertilization (IVF) in ewes during breeding season and seasonal anestrus". *Theriogenology* 56.1 (2001): 51-64.
26. Hasler John F. "Factors affecting frozen and fresh embryo transfer pregnancy rates in cattle". *Theriogenology* 56.9 (2001): 1401-1415.
27. Gonzalez-Bulnes A, *et al.* "Effects of FSH commercial preparation and follicular status on follicular growth and superovulatory response in Spanish Merino ewes". *Theriogenology* 54.7 (2000): 1055-1064.
28. Cognie Y. "State of the art in sheep-goat embryo transfer". *Theriogenology* 51.1 (1999): 105-116.
29. Nicholas FW. "Genetic improvement through reproductive technology". *Animal Reproduction Science* 42.1-4 (1996): 205-214.
30. Hasler JF, *et al.* "Production, freezing and transfer of bovine IVF embryos and subsequent calving results". *Theriogenology* 43.1 (1995): 141-152.
31. Choudhary KK, *et al.* "Advances in reproductive biotechnologies". *Veterinary world* 9.4 (2016): 388.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667