



## Obesity's Importance to *Equidae* and its Consequences

**Rodrigo Martins Ribeiro\* and Debora da Silva Freitas Ribeiro**

*Department of Veterinary Medicine, School of Veterinary Medicine, Centro Universitário de Mineiros, Mineiros, GO, Brazil*

**\*Corresponding Author:** Rodrigo Martins Ribeiro, Department of Veterinary Medicine, School of Veterinary Medicine, Centro Universitário de Mineiros, Mineiros, GO, Brazil.

**Received:** April 24, 2021

**Published:** April 28, 2021

© All rights are reserved by **Rodrigo Martins Ribeiro and Debora da Silva Freitas Ribeiro.**

Obesity is defined by World Health Organization as an abnormal or excessive accumulation of adipose tissue in the body [1]. According to Body Condition Score (BCS) developed by Henneke, et al. (1983), horses and ponies with BCS 7 can be considered overweight and BCS 8 or 9 are considered obese [2].

With the effect of being characterized an issue to equine creation getting worse by the fact that a lot of owners consider, until a certain point, obesity normal, acceptable or even desirable. In some competitions, the animal is rated by its physical characteristics, on these conditions, if the animal presents a certain degree of obesity, it can be considered as a physical beauty feature [3].

In developed countries, prevalence of obesity can be estimated around 19 to 40% in a equine's population [4,5]. To demonstrate how obesity is a common situation on this specie, a study assessing 300 adult animals, found out the rate of 19% obese animals (BCS between 8 and 9) and 32% presented overweight (BCS between 6.5 and 7.5) [6]. Was stressed in another study where a population of 319 animals randomly selected, it was found that 45% of these animals were considered obese or very obese, in South West region of Goias, located in Brazil was found a total 12.16% of equines with BCS above 7 [7,8].

Obesity induction occurs with supply of diets in grains and forage (grass and hay) with high contents of non-structural carbohydrates, becoming therefore a consequence of over-supply, providing excess of metabolic exigences to its physical activity level [3,9].

When adipocytes reach its maximum storage capacity, changes such as energy inefficiency, inflammatory processes and cell stress begins. Even hypoxia cases can happen in adipose tissues. This occurs due to excess of fat accumulation inside adipose tissues, that therefore, expand, compromising oxygen transport through compression of adjacent capillaries. Along this situation, these endothelium capillaries may not respond to nitric oxide, decreasing its vasodilation. These adipocytes hypoxia cases will compromise mitochondrial action and therefore end up stimulating inflammatory process, releasing cytokines and macrophages chemotactic proteins [10].

It's important to highlight that adipose tissue can no longer be considered as an organ that just stores energy, but also as an endocrine organ with paracrine and autocrine action. This tissue is responsible for producing hormones such as adipokines, that are released from adipocytes and other resident cells on adipose tissue. Adipocytes can excrete leptin, resistin, adiponectin, vistafin and apelin. Other than that, releases inflammatory cytokines such as a tumor necrosis factor-alpha (TNF- $\alpha$ ), interleukins 1 (IL-1) and 6 (IL-6), macrophages chemotactic protein (MCP-1), macrophage inhibition factor (MIF), besides inflammatory proteins such as haptoglobin, inhibitor of plasminogen activator 1(PAI-1) serum amyloid protein A (SAA) [11-15].

In response to obesity, animals present cases of exercise intolerance, thermoregulatory inefficiency, abnormal reproductive development, and probability increase in mesenteric lipomas development [16-19].

Thereby, the issue about overweight in horses must be remembered by veterinarians and professionals that work with equine production, being necessary heightened attention about its harmful effects and consequences on these animals' life quality.

## Bibliography

1. Geor R J. "Metabolic Predispositions to Laminitis in Horses and Ponies: Obesity, Insulin Resistance and Metabolic Syndromes". *Journal of Equine Veterinary Science* 28.12 (2008): 753-759.
2. Johnson PJ, et al. "Laminitis and the equine metabolic syndrome". *Veterinary Clinics of North America: Equine Practice* 26 (2010): 239-255.
3. Wild S H and BYRNE C D. "ABC of obesity: risk factors for diabetes and coronary heart disease". *British Medical Journal* 333 (2006): 1009-1011.
4. Wyse C A., et al. "Prevalence of obesity in riding horses in Scotland". *Veterinary record* 162 (2008): 590-591.
5. Thatcher C D., et al. "Prevalence of over conditioning in mature horses in southwest Virginia during the summer". *Journal of Veterinary Internal Medicine* 26 (2012): 1413-1418.
6. Allan F J., et al. "A cross-sectional study of risk factors for obesity in cats in New Zealand". *Preventive Veterinary Medicine* 46.3 (2000): 183-196.
7. Robertson ID. "The association of exercise, diet and other factors with owner-perceived obesity in privately owned dogs from metropolitan PERTH, W. A". *Preventive Veterinary Medicine* 58.1-2 (2003): 75-83.
8. Ribeiro R M., et al. "Avaliação do escore corporal e acúmulo de adiposidades regionais dos equinos localizados no sudoeste goiano". In: XVI CONGRESSO INTERNACIONAL VETERINÁRIO MONTENEGRO, 2020, Santa Maria. Anais XVI CONGRESSO INTERNACIONAL VETERINÁRIO MONTENEGRO (2020): 76-77.
9. Schott HC., et al. "The Michigan Cushing's project". In: Proceedings of the 47th Annual Convention of the American Association of Equine Practitioners (2001): 22-24.
10. Goossens G H. "The role of adipose tissue dysfunction in the pathogenesis of obesity-related insulin resistance". *Physiology Behavior* 94 (2008): 206-218.
11. Gerhardt CC., et al. "Chemokines control fat accumulation and leptin secretion by cultured human adipocytes". *Molecular and Cellular Endocrinology* 175 (2001): 81-92.
12. Sartipy P., et al. "Monocyte chemoattractant protein 1 in obesity and insulin resistance". *Proceedings of the National Academy of Sciences of the United States of America* 100 (2003): 7265-7270.
13. Rajala MW and SCHERER PE. "Minireview: the adipocyte—at the crossroads of energy homeostasis, inflammation, and atherosclerosis". *Endocrinology* 144 (2003): 3765-3773.
14. Trayhurn P and WOOD IS. "Adipokines: inflammation and the pleiotropic role of white adipose tissue". *British Journal of Nutrition* 92 (2004): 347-355.
15. Rasouli N and KERN PA. "Adipocytokines and the metabolic complications of obesity". *The Journal of Clinical Endocrinology and Metabolism* 93 (2008): 564-573.
16. HENNEKE DR., et al. "Body condition during pregnancy and lactation and reproductive efficiency of mares". *Theriogenology* 21 (1983): 897-909.
17. Cymbalukm NF and CHRISTISON GI. "Environmental effects on thermoregulation and nutrition of horses". *Veterinary Clinics of North America: Equine Practice* 6.2 (1990): 355-372.
18. Garlinghouse SE and Burrill MJ. "Relationship of body condition score to completion rate during 160 km endurance races". *Equine Veterinary Journal* (1999): 591-595.

19. Garcia-Seco E., *et al.* "Prevalence and risk factors associated with outcome of surgical removal of pedunculated lipomas in horses: 102 cases (1987-2002)". *Journal of the American Veterinary Medical Association* 226.9 (2005): 1529-1537.

**Assets from publication with us**

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

**Website:** [www.actascientific.com/](http://www.actascientific.com/)

**Submit Article:** [www.actascientific.com/submission.php](http://www.actascientific.com/submission.php)

**Email us:** [editor@actascientific.com](mailto:editor@actascientific.com)

**Contact us:** +91 9182824667