



Figure 2: Mountain Roof Top Forests (MRTFs) in Western Ghats.

Fresh water streams, canals and pools are found along the course of the Keeriparai Forest Range. As rubber cultivation is predominant in this area, the monoculture exhausted the soil and needed a lot of pesticides and fertilizers. This would have caused soil and water pollution that gives negative consequences to the natural environment [6]. In this eco system, because of their size and interesting behavior, dragonflies and damselflies are considered to be the most charismatic of all aquatic insects.



Figure 3: Tropical Evergreen Forests.

In the invertebrate world of Odonates, *Neurobasis chineseis* is always considered an ecological indicator to study the eco system. The male will have abdomen of size 45-50 mm, hind wings of 32-38 mm with eyes blackish brown above and bluish white below [9]. The legs are dark bronze with white outer stripe, rounded wings at tips with forewings transparent, tinted with pale yellowish green with emerald green venation.

The hind wings are opaque, two thirds iridescent green or peacock blue. Apical half is blackish brown with violet reflections and green iridescent veins. Underside of hind wings will be uniformly blackish brown with dull golden reflections. The abdomen will be iridescent green above and on side and underside will be black. The 9th and 10th segments are whitish in the abdomen.



Figure 4: Male *Neurobasis chineseis*.

The female will have abdomen of length 44-50mm, hind wings of length 36-40 mm. Eyes will be brownish above and yellowish below [9]. The thorax and legs are similar to that of males. The wings are transparent and amber colored. All wings have a round creamy white central opaque spot on the edge of the wing. Wing spot is absent in forewings and creamy white in hind wings. The abdomen is dull iridescent green above and black below. Green metallic stripe on sides in all segments bordered with black.



Figure 5: Female *Neurobasis chineseis*.

The habitat of *Neurobasis chineseis* is found between 500 - 1200 m altitude. It perches on emergent boulders and fallen logs in streams. The bright green dorsal iridescence of the hind wings of *Neurobasis chineseis* males, very rare in Odonata, is known to play a significant role in their courtship behavior [2]. Female lay eggs on submerged decaying logs in streams during southwest monsoon. The flight season is from May to November.

Materials and Methods

Study area

The study was conducted in the Keeriparai Forest Range 29 km², a part of Agasthiyamalai 8.23 Latitude and 77.50 Longitude located in the southern end of Western Ghats. Geologically, the rocks are granitoid gneiss and the terrain is undulating [1]. The



Figure 6: Keeriparai Forest Range.

mean annual ambient temperature in the evergreen forest is 22.5°C (range 15–30°C). The mean annual rainfall is about 3,000 mm from the southwest monsoon (May-August) and northeast monsoons (October-December) [7]. The type of vegetation varies along the elevation gradient. The foothills harbor rubber plantations up to an altitude of 200 m. Dry teak, composed of stunted, thin-poled teak (*Tectona grandis*), and rubber plantations occurs between 200 and 850 m, wet evergreen forest from 920 to 1,500 m, beyond which high-altitude Mountain Roof Top Forests (MRTFs) occupy.



Figure 7: Habitat of *Neurobasis chineseis*.

Data collection

Species survey

The abundance of *Neurobasis chineseis* dragonfly species is directly proportional to the vascular plants they feed in freshwater habitat due to unpolluted water streams, emphasis the ecological indicator of the richness of the forest eco system.

The observation of *Neurobasis chineseis* was carried out between 9:00 and 15:00 hrs., which correspond to the period of greatest activity [3]. A span of 29 kms was surveyed from December 2021

to March 2022. In order to determine the abundance and species richness, the *Neurobasis chineseis* were observed along streams, canals and pools as transect walk around 29 sq. kms within the forest range at 100 to 800 meter altitude.

Data analysis

To explore the abundance of *Neurobasis chineseis*, environment analysis of water quality and richness of plants along the freshwater systems were considered as main parameters. As unpolluted water is the main source for the growth of vascular plants on which *Neurobasis chineseis* feeds. Hence for water analysis, simple parameters of pH, COD and BOD were analyzed to determine the water quality. For vegetation, the types of plants found along the streams and also the moss and likens varieties on the rocks in streams were analyzed to find the abundance of *Neurobasis chineseis*. These parameters were considered to validate the hypothesis of *Neurobasis chineseis* as an Ecological Indicator to assess the richness of the eco system in Keeriparai Forest Range of Kanyakumari District.

Observations and Analysis

The water samples were tested for the pH value to assess the acidity/alkaline nature of the stream water at different altitudes. The details of the analysis are given below.

Height (Meters)	pH
100	7.1
300	7.0
500	7.0
700	7.0
900	7.0

Table 1

It is clear from the table that there is no change in the pH value, as the water is not polluted by any fertilizers used for enriching the soil for plantation of rubber crops. Also the water bodies are not used as a dumping ground for plastics or other household wastes.

The Biochemical Oxygen Demand (BOD) levels of stream water is being tested and the details are given below.

Height (Meters)	BOD (mg/l)
100	0.8
300	0.7
500	0.7
700	0.7
900	0.7

Table 2

From the table, it is clear that no domestic waste is being discharged in the river stream. Even though there are workers involved in estate plantations they are aware of protecting the eco system.

The Chemical Oxygen Demand (COD) level of stream water at different altitudes were analyzed and the details are given below.

Height (Meters)	COD (mg/l)
100	18.42
300	17.71
500	17.71
700	17.71
900	17.71

Table 3

Commercial crops like rubber and clove are cultivated in the forest range. As it is monoculture, there is a threat of fertilizers to be used that will pollute the water system. Also cleaning of rubber containers will also change the COD level of water. But from our analysis that there is no pollution of water bodies.

The Western Ghats harbor about 320 species of ferns and fern allies with more species diversity in the southern part [8]. The major families of pteridophytes found in the Western Ghats are Aspleniaceae, Polypodiaceae, Thelypteridaceae, Selaginellaceae, Pteridaceae, etc. Whereas on the generic level, maximum diversity is observed in the genus *Asplenium*, *Selaginella*, *Pteris*, *Athyrium*, *Diplazium*, etc. In the Keeriparai Forest Range, the habitat of the pteridophytes consists of microclimatic conditions with special preference for moist and shady places. As ferns need moist environment, along the course of stream water, plenty of ferns are observed near the banks and also on rock crevices.

Due to frequent rainfall in this region (3000 mm on an average), there is water flow in the stream throughout the year which is the favorable condition for the survival of fern varieties. Hence the growth of ferns along the streams provide good source of food for *Neurobasis chinesis* throughout the year which is unique in this eco system.

Results and Discussions

From our study, the occurrence *Neurobasis chinesis* in the altitude of 700 to 900 meters in the Keeriparai range is strong evidence that the freshwater stream is not polluted. Cash crops like rubber and clove are cultivated that need fertilizers to enrich the soil. Our analysis of water samples revealed that the stream water is not polluted. Our observations and discussions revealed that the farmers use natural biomass to make the soil fertile. The forest dwellers use the stream water for drinking purpose and hence they avoid washing clothes or discharging domestic wastes in the stream water. No chemical pollution is found in the stream as the farmers wash rubber sheets and latex containers away from the stream water. As Keeriparai Range comes under Protected Area, tourism is not allowed and there is no evidence of plastic wastes or debris near the stream.

Conclusion

Hence our study reveals that the farmers and forest dwellers are aware of managing the eco system with their traditional knowledge, as *Neurobasis chinesis* is found in this forest range that serves as an Ecological Indicator for the richness of Western Ghats in Kanyakumari District.

Bibliography

1. Amarnath Giriraj., *et al.* "A method for assessing evergreen habitats using Phyto diversity and geospatial techniques in tropical rain forests of Southern Western Ghats (India)". *Ecological Research* 24.4 (2009): 749-760.
2. Andre Gunther and Dagmar Hilfert-Reuppel. "Reproductive Behavior and the system of signaling in *Neurobasis Chinesis* (Odonata Calopterygidae - a kinematic analysis)". *International Journal of Odonatology* (2014): 31-52.
3. Corbet PS. "Dragonflies: Behaviour and Ecology of Odonata". *International Journal of Freshwater Entomology* 23.1 (1999): 83.

4. R Gopalan and AN Henry. "Endemic Plants of India- Endemics of Agasthiyamalai Hills". Bishen Singh Ma- hendrapal Singh publications, Dehradun (2000).
5. D Narasimham and Sheeba J Erwin. "Database of Agasthiyamalai Biosphere Reserve". Report submitted to Department of Forest, Tamil Nadu (2017).
6. Bijaya Kumar Panda and Sukanta Sarkat. "Environment Impact of Rubber Plantation: Ecological Vs. Economical Perspective". *Asian Journal of Microbiology, Biotechnology and Environmental Sciences* 22.4 (2020): 657-661.
7. Pascal JP. "Wet Evergreen Forest of the Western Ghats of India. Ecology, Structure, floristic composition and succession". Institut Francais De Pondicherry (1988): 345.
8. Sumesh Dudani., *et al.* "Pteridophytes of Western Ghats". *Biodiversity Documentation and Taxonomy Published by Narendra Publishing House* (2009).
9. P Vukusic., *et al.* "Remarkable iridescence in the hindwings of the damselfly *Neurobasis chinensis chinensis*". *Proceedings of the Royal Society B* 271.1539 (2003): 595-601.