



Bridging the Gap: Enhancing Knowledge, Attitudes, and Perceptions of Polycystic Ovarian Syndrome (PCOS)

Sawera Haider*, Syed Muhammad Abbas and Ashhad Rabeegh

Department of Medicine, Dow University of Health Sciences, Karachi, Pakistan

*Corresponding Author: Sawera Haider, Department of Medicine, Dow University of Health Sciences, Karachi, Pakistan.

Received: August 05, 2024

Published: August 12, 2024

© All rights are reserved by Sawera Haider., et al.

I am writing to address the crucial topic of knowledge, attitudes, and perceptions about Polycystic Ovarian Syndrome (PCOS). PCOS is a common endocrine disorder affecting women of reproductive age, characterised by symptoms such as irregular menstrual cycles, hyperandrogenism, and polycystic ovaries. Despite its prevalence, there remains a significant gap in public knowledge and understanding of this condition, which can hinder early diagnosis and effective management.

Firstly, it is essential to recognize that public awareness about PCOS is relatively low. A study by Balen., et al. found that only a small percentage of women are aware of PCOS and its symptoms, which leads to delays in seeking medical help [1]. This lack of awareness extends to healthcare providers as well, with many general practitioners not fully versed in the diagnostic criteria and treatment options for PCOS [2]. The limited exposure to PCOS during medical training further exacerbates this issue, resulting in misdiagnosis or delayed diagnosis.

The attitudes towards PCOS can also significantly impact the quality of life for those affected. Women with PCOS often report feeling stigmatized and misunderstood, both by their peers and by healthcare professionals [3]. This stigma can lead to psychological distress, including anxiety and depression, which are already more prevalent in women with PCOS compared to the general population [4]. The physical symptoms of PCOS, such as hirsutism, acne, and weight gain, can also contribute to poor self-esteem and body image issues. Addressing these psychosocial aspects is crucial in providing holistic care to women with PCOS.

Moreover, the perceptions surrounding the management of PCOS are often flawed. There is a common misconception that PCOS can be completely cured, leading to unrealistic expectations and disappointment when managing the chronic nature of the condition [5]. Education campaigns that emphasise the importance of lifestyle modifications, such as diet and exercise, alongside medical treatments, are crucial for improving long-term outcomes [6].

It is important for patients to understand that while PCOS cannot be cured, its symptoms can be effectively managed with a combination of lifestyle changes and medical interventions. Regular follow-ups with healthcare providers are essential to monitor and adjust treatment plans as needed.

Efforts to bridge the knowledge gap must involve comprehensive educational programs aimed at both the public and healthcare providers. Resources such as the Rotterdam criteria, which offer a standardized diagnostic framework, should be more widely disseminated and utilised [7]. Additionally, support groups and online communities can play a vital role in providing information and emotional support to those affected by PCOS [8]. These platforms can also help in sharing personal experiences and coping strategies, fostering a sense of community and reducing feelings of isolation among women with PCOS.

Research is another critical area that requires attention. There is a need for more extensive studies to understand the etiology of PCOS better and develop more effective treatment strategies. The heterogeneity of PCOS symptoms and their impact on different ethnic groups highlight the importance of personalized medicine in managing this condition. Collaborative research efforts can pave the way for new discoveries and innovations in PCOS treatment, ultimately improving the quality of life for women with this disorder.

Healthcare policies should also be re-evaluated to provide better support for women with PCOS. Insurance coverage for diagnostic tests and treatments, including fertility treatments, should be more inclusive to reduce the financial burden on patients. Additionally, integrating PCOS education into school health programs can help in early identification and management, empowering young women to seek medical advice promptly.

In conclusion, increasing awareness, improving attitudes, and correcting misconceptions about PCOS are critical steps towards better management and support for women with this condition.

Collaborative efforts between healthcare providers, educators, and the media are essential in promoting a more informed and supportive environment for those impacted by PCOS. By addressing these issues comprehensively, we can ensure that women with PCOS receive the care and support they need to lead healthy and fulfilling lives.

Bibliography

1. Balen A H., *et al.* "Polycystic ovary syndrome: the spectrum of the disorder in 1741 patients". *Human Reproduction* 10.8 (1995): 2107-2111.
2. Teede H J., *et al.* "Guidelines for the assessment and management of polycystic ovary syndrome". *Medical Journal of Australia* 195.6 (2011): S65-S112.
3. Himelein M J and Thatcher S S. "Polycystic ovary syndrome and mental health: A review". *Obstetrical and Gynecological Survey* 61.11 (2006): 723-732.
4. Dokras A., *et al.* "Screening women with polycystic ovary syndrome for metabolic syndrome". *Obstetrics and Gynecology* 106.1 (2005): 131-137.
5. Azziz, R., *et al.* "Polycystic ovary syndrome". *Nature Reviews Disease Primers* 2 (2016): 16057.
6. Moran LJ., *et al.* "Diet and exercise in polycystic ovary syndrome: lifestyle management for insulin resistance related infertility". *Seminars in Reproductive Medicine* 26.1 (2008): 58-65.
7. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. "Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome". *Fertility and Sterility* 81.1 (2004): 19-25.
8. Kitzinger C and Willmott, J. "'The thief of womanhood': Women's experience of polycystic ovarian syndrome". *Social Science and Medicine* 54.3 (2002): 349-361.