



Safeguarding Health: Unveiling the Essentials of STI Diagnosis and Treatment

Sawera Haider*, Ateesh Kumar, Ravesh Kumar and Anoshia Ali

Department of Internal Medicine, Dow University of Health Sciences, Pakistan

*Corresponding Author: Sawera Haider, Department of Internal Medicine, Dow University of Health Sciences, Pakistan.

Received: August 13, 2024

Published: August 30, 2024

© All rights are reserved by Sawera Haider., et al.

Dear Editor,

I am writing to express my growing concern over the rising rates of sexually transmitted infections (STIs) in our society, particularly among the younger population. The prevalence of unsafe sexual practices has significantly contributed to this public health crisis, which, if not addressed promptly, could have far-reaching consequences for both individual and community health. It is imperative that we focus on increasing awareness about STIs, promoting safe sexual practices, and ensuring that young people are equipped with the knowledge and resources they need to protect themselves.

The surge in STIs is a global issue, but it is especially troubling among young people aged 15-24, who account for a significant proportion of new infections. According to the World Health Organization (WHO), more than 1 million STIs are acquired every day worldwide, with young people disproportionately affected [1]. This age group is often at higher risk due to a combination of factors, including lack of knowledge about sexual health, inconsistent use of protection, and engagement in high-risk behaviours such as having multiple sexual partners [2].

One of the primary drivers of this increase in STIs is the widespread lack of comprehensive sexual education. Many young people are not adequately informed about the importance of safe sex, the correct use of condoms, and the serious health consequences of untreated STIs. Studies have shown that inadequate sexual education contributes to risky behaviours and a higher likelihood of contracting STIs [3]. This gap in education leaves many young individuals unprepared to make informed decisions about their sexual health, putting them at greater risk of infection.

The rise of casual sexual encounters, often facilitated by social media and dating apps, has further exacerbated the problem. These platforms have made it easier for individuals to engage in sexual activities with multiple partners, often without taking the necessary precautions to prevent STIs [4]. This trend underscores the urgent need for targeted educational campaigns that address

the realities of modern sexual behaviour and emphasize the importance of safe practices.

Moreover, the stigma associated with STIs continues to be a significant barrier to effective prevention and treatment. Many young people feel ashamed or embarrassed to seek information, get tested, or discuss sexual health with healthcare providers [5]. This reluctance to engage in open conversations about sexual health leads to a lack of awareness and delays in seeking treatment, which can result in severe health complications and further spread of infections.

Public awareness and education are critical components in the fight against STIs. Comprehensive sexual education should be implemented in schools and communities, with a focus on teaching young people about safe sexual practices, the correct use of condoms, and the importance of regular STI testing [6]. Education should also address the emotional and social aspects of sexual health, helping young people navigate relationships and make informed decisions about their sexual behaviour.

Condom use remains one of the most effective methods of preventing STIs, yet many young people lack the knowledge or motivation to use them consistently [7]. Public health campaigns should prioritize educating young people on the correct and consistent use of condoms and other forms of protection. Additionally, efforts should be made to ensure that condoms are widely available, affordable, and accessible to all, particularly in areas with high rates of STIs [8].

Regular screening and early diagnosis are also essential in reducing the spread of STIs. Many STIs, such as chlamydia and gonorrhoea, are often asymptomatic, meaning that individuals may be unaware they are infected and unintentionally transmit the infection to others [9]. By promoting regular STI testing as a standard part of healthcare for sexually active individuals, we can identify and treat infections early, preventing further transmission and reducing the overall burden of STIs in the community.

In addition to education and prevention, it is crucial to address the issue of stigma that surrounds STIs. Stigma not only discourages individuals from seeking care but also perpetuates misinformation and fear. We must work to create an environment where discussing sexual health is normalized, and where individuals feel comfortable seeking the care and information they need without fear of judgement. This can be achieved through public awareness campaigns, community outreach, and by fostering open dialogues about sexual health in both public and private settings [10].

Furthermore, healthcare providers play a vital role in combating the STI epidemic. They must be equipped with the knowledge and resources to provide accurate information, offer confidential and non-judgmental care, and encourage regular screening for STIs. Providing young people with access to affordable and confidential sexual health services is essential in ensuring that they receive the care they need in a timely manner [11].

In conclusion, the rising rates of STIs among the younger population are a direct consequence of unsafe sexual practices and a lack of comprehensive knowledge about sexual health. To address this public health crisis, we must focus on enhancing sexual education, promoting safer sex practices, and reducing the stigma associated with STIs. By empowering young people with the knowledge and resources they need to make informed decisions about their sexual health, we can significantly reduce the incidence of STIs and protect the health of future generations.

Bibliography

1. World Health Organization. Sexually transmitted infections (STIs) (2021).
2. Centers for Disease Control and Prevention. Sexually Transmitted Infections Treatment Guidelines (2022).
3. Lindberg L D., *et al.* "Changes in adolescents' receipt of sex education, 2006–2013". *Journal of Adolescent Health* 58.6 (2016): 621-627.
4. Landovitz R J., *et al.* "Prevalence of HIV, sexually transmitted infections, and sexual risk behavior among young men who have sex with men: The effect of recent sex work". *Archives of Sexual Behavior* 42.4 (2013): 678-689.
5. Fenton KA., *et al.* "Measuring sexual behavior: methodological challenges in survey research". *Sexually Transmitted Infections* 77.2 (2001): 84-92.
6. Workowski K A and Bolan G A. "Sexually transmitted diseases treatment guidelines, 2015". *MMWR Recommendations and Reports* 64 (RR-03) (2015): 1-137.
7. Reece M., *et al.* "Condom use rates in a national probability sample of males and females ages 14 to 94 in the United States". *Journal of Sexual Medicine* 7.5 (2010): 266-276.
8. Kirby D B. "Emerging answers 2007: Research findings on programs to reduce teen pregnancy and sexually transmitted diseases". National Campaign to Prevent Teen and Unplanned Pregnancy.
9. Hook EW and Handsfield H H. "Gonococcal infections in the adult". *Sexually Transmitted Diseases* 3.9 (2008): 873-884.
10. Brown AE., *et al.* "Sexual health and relationships in the UK: National findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3)". *The Lancet* 382.9907 (2008): 1845-1855.
11. Guttmacher Institute. "Adolescents' Need for and Use of Sexual Health Services in the United States" (2019).